

Datasheet: MCA342GA

Description:	MOUSE ANTI RAT CD163
Specificity:	CD163
Other names:	ED2
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	ED2
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			
Immunohistology - Frozen	▪			1/50 - 1/100
Immunohistology - Paraffin (1)	▪			
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting	▪			
Immunofluorescence	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) This product requires protein digestion pre-treatment of paraffin sections e.g. trypsin or pronase

Target Species	Rat
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline

Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 0.5 mg/ml
Immunogen	Rat Spleen cell homogenate.
RRID	AB_2074558
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the SP2/0-Ag 14 mouse myeloma cell line.
Specificity	<p>Mouse anti Rat CD163, clone ED2 recognizes the rat ED2 cell surface glycoprotein (Dijkstra et al. 1985). A 175 kDa molecule also known as rat CD163, a member of the group B scavenger receptor cysteine-rich (SRCR) family and an erythroblast adhesion receptor (Fabriek et al. 2007).</p> <p>Mouse anti rat CD163, clone ED2 was shown to detect approximately 50% of peritoneal macrophages, a subset of splenic macrophages, and most tissue macrophages. However, no staining was observed in monocytes or alveolar macrophages (Dijkstra et al. 1985, Beelen et al. 1987). In freshly isolated bone marrow, expression of CD163 was limited to mature macrophages only (Barbe et al. 1990).</p> <p>Clone ED2 may be used in immunohistology using antigen retrieval, and has also been described reacting with paraffin-embedded material following PLP fixation (Periodate-lysine-paraformaldehyde), see Whiteland et al.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul
Histology Positive Control Tissue	Liver
References	<ol style="list-style-type: none"> Dijkstra, C.D. <i>et al.</i> (1985) The heterogeneity of mononuclear phagocytes in lymphoid organs: distinct macrophage subpopulations in the rat recognized by monoclonal antibodies ED1, ED2 and ED3. Immunology. 54 (3): 589-99. Beelen, R.H.J. <i>et al.</i> (1987) Monoclonal antibodies ED1, ED2 and ED3 against rat macrophages: Expression of recognized antigens in different stages of differentiation. Transplant Proc. 3: 3166-70. Barbe, E. <i>et al.</i> (1990) Characterization and expression of the antigen present on resident rat macrophages recognized by monoclonal antibody ED2. Immunobiol. 182: 88-99. Dijkstra, C.D. & Damoiseaux, J.G. (1993) Macrophage heterogeneity established by immunocytochemistry. Prog Histochem Cytochem. 27 (2): 1-65. Whiteland, J.L. <i>et al.</i> (1995) Immunohistochemical detection of T-cell subsets and other leukocytes in paraffin-embedded rat and mouse tissues with monoclonal antibodies. J Histochem Cytochem. 43 (3): 313-20.

6. Polfliet, M.M.J. *et al.* (2002) Identification of the rat mature macrophage antigen ED2 as CD163: Regulation by glucocorticoids and role in the production of proinflammatory mediators. PhD Thesis. Vrije University, Amsterdam.
7. Deng, X. *et al.* (2005) Chronic alcohol consumption accelerates fibrosis in response to cerulein-induced pancreatitis in rats. [Am J Pathol. 166 \(1\): 93-106.](#)
8. Piquet-Pellorce, I. *et al.* (2005) Identification of the leukemia inhibitory factor cell targets within the rat testis. [Biol Reprod. 72: 602-11.](#)
9. Fujita, E. *et al.* (2010) Statin attenuates experimental anti-glomerular basement membrane glomerulonephritis together with the augmentation of alternatively activated macrophages. [Am J Pathol. 177 \(3\): 1143-54.](#)
10. Schwartzkopff, J. *et al.* (2010) NK cell depletion delays corneal allograft rejection in baby rats. [Mol Vis. 16: 1928-35.](#)
11. Baker, S.C. *et al.* (2011) Cellular integration and vascularisation promoted by a resorbable, particulate-leached, cross-linked poly(ϵ -caprolactone) scaffold. [Macromol Biosci. 11 \(5\): 618-27.](#)
12. Bedi, A. *et al.* (2010) Effect of early and delayed mechanical loading on tendon-to-bone healing after anterior cruciate ligament reconstruction. [J Bone Joint Surg Am. 92: 2387-401.](#)
13. Banerjee, S. *et al.* (2003) Development of organised conjunctival leucocyte aggregates after corneal transplantation in rats. [Br J Ophthalmol. 87: 1515-22.](#)
14. Moghaddami, M. *et al.* (2005) MHC class II compartment, endocytosis and phagocytic activity of macrophages and putative dendritic cells isolated from normal tissues rich in synovium. [Int Immunol. 17: 1117-30.](#)
15. Wojcik, M. *et al.* (2012) Immunodetection of cyclooxygenase-2 (COX-2) is restricted to tissue macrophages in normal rat liver and to recruited mononuclear phagocytes in liver injury and cholangiocarcinoma. [Histochem Cell Biol. 137: 217-33.](#)
16. Kawakami, A.P. *et al.* (2011) Inflammatory Process Modulation by Homeopathic Arnica montana 6CH: The Role of Individual Variation. [Evid Based Complement Alternat Med. 2011: 917541.](#)
17. Kajita, M. *et al.* (2011) iNOS expression in vascular resident macrophages contributes to circulatory dysfunction of splanchnic vascular smooth muscle contractions in portal hypertensive rats. [Am J Physiol Heart Circ Physiol. 300: H1021-31.](#)
18. Jiao, K. *et al.* (2013) The Identification of CD163 Expressing Phagocytic Chondrocytes in Joint Cartilage and Its Novel Scavenger Role in Cartilage Degradation. [PLoS One. 8\(1\):e53312.](#)
19. Fujii, Y. *et al.* (2013) Effect of enzymatically modified isoquercitrin on preneoplastic liver cell lesions induced by thioacetamide promotion in a two-stage hepatocarcinogenesis model using rats. [Toxicology. 305: 30-40.](#)
20. Lobato-Pascual, A. *et al.* (2013) Rat macrophage C-type lectin is an activating receptor expressed by phagocytic cells. [PLoS One. 8: e57406.](#)
21. Fabriek, B.O. *et al.* (2007) The macrophage CD163 surface glycoprotein is an erythroblast adhesion receptor. [Blood. 109 \(12\): 5223-9.](#)
22. Park, E.S. *et al.* (2014) Establishment of a rat model for canine necrotizing meningoencephalitis (NME). [Vet Pathol. 51 \(6\): 1151-64.](#)
23. Keitel, V. *et al.* (2008) Expression and function of the bile acid receptor TGR5 in Kupffer cells. [Biochem Biophys Res Commun. 372: 78-84.](#)
24. Hozumi, Y. *et al.* (2015) Expression and localization of the diacylglycerol kinase family

- and of phosphoinositide signaling molecules in adrenal gland. [Cell Tissue Res. 362 \(2\): 295-305.](#)
25. Fernandez-Bustamante, A. *et al.* (2015) Brief Glutamine Pretreatment Increases Alveolar Macrophage CD163/Heme Oxygenase-1/p38-MAPK Dephosphorylation Pathway and Decreases Capillary Damage but Not Neutrophil Recruitment in IL-1/LPS-Insufflated Rats. [PLoS One. 10 \(7\): e0130764.](#)
26. Adamo, H.H. *et al.* (2015) Adaptive (TINT) Changes in the Tumor Bearing Organ Are Related to Prostate Tumor Size and Aggressiveness. [PLoS One. 10 \(11\): e0141601.](#)
27. Santos, L. *et al.* (2016) *In vitro* and *in vivo* assessment of magnetically actuated biomaterials and prospects in tendon healing. [Nanomedicine \(Lond\). 11 \(9\): 1107-22.](#)
28. Ibarra V *et al.* (2016) Evaluation of the Tissue Response to Alginate Encapsulated Islets in an Omentum Pouch Model. [J Biomed Mater Res A. May 3. \[Epub ahead of print\]](#)
29. Tentillier N *et al.* (2016) Anti-Inflammatory Modulation of Microglia via CD163-Targeted Glucocorticoids Protects Dopaminergic Neurons in the 6-OHDA Parkinson's Disease Model. [J Neurosci. 36 \(36\): 9375-90.](#)
30. Zakrzewicz, A. *et al.* (2015) Monocytic Tissue Transglutaminase in a Rat Model for Reversible Acute Rejection and Chronic Renal Allograft Injury. [Mediators Inflamm. 2015: 429653.](#)
31. Rave-Fränk M *et al.* (2013) Rat model of fractionated (2 Gy/day) 60 Gy irradiation of the liver: long-term effects. [Radiat Environ Biophys. 52 \(3\): 321-38.](#)
32. Wang, M. *et al.* (2017) Characterization of the Micro-Environment of the Testis that Shapes the Phenotype and Function of Testicular Macrophages. [J Immunol. May 1. pii: 1700162. \[Epub ahead of print\]](#)
33. Stavenuiter, A.W. *et al.* (2015) Protective Effects of Paricalcitol on Peritoneal Remodeling during Peritoneal Dialysis. [Biomed Res Int. 2015: 468574.](#)
34. Hawkins, K.E. *et al.* (2017) Targeting resolution of neuroinflammation after ischemic stroke with a lipoxin A₄ analog: Protective mechanisms and long-term effects on neurological recovery. [Brain Behav. 7 \(5\): e00688.](#)
35. Almahrog, A.J. *et al.* (2016) *In vivo* association of immunophenotyped macrophages expressing CD163 with PDGF- β in gingival overgrowth-induced by three different categories of medications. [J Oral Biol Craniofac Res. 6 \(1\): 10-7.](#)
36. Han, T.T. *et al.* (2015) Adipose-derived stromal cells mediate *in vivo* adipogenesis, angiogenesis and inflammation in decellularized adipose tissue bioscaffolds. [Biomaterials. 72: 125-37.](#)
37. Pannell, M. *et al.* (2016) Adoptive transfer of M2 macrophages reduces neuropathic pain via opioid peptides. [J Neuroinflammation. 13 \(1\): 262.](#)
38. Chang, J.C. *et al.* (2019) Early Immune Response to Acute Gastric Fluid Aspiration in a Rat Model of Lung Transplantation. [Exp Clin Transplant. 17 \(1\): 84-92.](#)
39. Bloomer, S.A. *et al.* (2019) Aging results in accumulation of M1 and M2 hepatic macrophages and a differential response to gadolinium chloride [Histochem Cell Biol. 25 Oct \[Epub ahead of print\].](#)
40. Azam, M. *et al.* (2019) Addition of 2-deoxy-d-ribose to clinically used alginate dressings stimulates angiogenesis and accelerates wound healing in diabetic rats. [J Biomater Appl. 34 \(4\): 463-475.](#)
41. Bhandari, S. *et al.* (2020) Transcriptome and proteome profiling reveal complementary scavenger and immune features of rat liver sinusoidal endothelial cells and liver macrophages. [BMC Mol Cell Biol. 21 \(1\): 85.](#)

42. Solár, P. *et al.* (2020) Subarachnoid Hemorrhage Increases Level of Heme Oxygenase-1 and Biliverdin Reductase in the Choroid Plexus. [Front Cell Neurosci. 14: 593305.](#)
43. Nishida, Y. *et al.* (2020) Intra-Articular Injection of Stromal Cell-Derived Factor 1 α Promotes Meniscal Healing via Macrophage and Mesenchymal Stem Cell Accumulation in a Rat Meniscal Defect Model. [Int J Mol Sci. 21\(15\):5454.](#)
44. Bloomer, S.A. *et al.* (2020) Aging results in accumulation of M1 and M2 hepatic macrophages and a differential response to gadolinium chloride. [Histochem Cell Biol. 153 \(1\): 37-48.](#)
45. Kuramochi, M. *et al.* (2021) Involvement of neutrophils in rat livers by low-dose thioacetamide administration. [J Vet Med Sci. Jan 20 \[Epub ahead of print\].](#)
46. Broekaart, D.W.M. *et al.* (2020) Increased matrix metalloproteinases expression in tuberous sclerosis complex: modulation by microRNA 146a and 147b *in vitro*.. [Neuropathol Appl Neurobiol. 46 \(2\): 142-159.](#)
47. Duchesne, E. *et al.* (2013) Mast cells can regulate skeletal muscle cell proliferation by multiple mechanisms. [Muscle Nerve. 48 \(3\): 403-14.](#)
48. Li, R. *et al.* (2020) Bone marrow mesenchymal stem cell-derived exosomal microRNA-124-3p attenuates neurological damage in spinal cord ischemia-reperfusion injury by downregulating Ern1 and promoting M2 macrophage polarization. [Arthritis Res Ther. 22 \(1\): 75.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Rabbit Anti Mouse IgG (STAR8...)	DyLight®800
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (STAR70...)	FITC
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP

Rabbit Anti Mouse IgG (STAR13...)

[HRP](#)

Goat Anti Mouse IgG (H/L) (STAR117...)

[Alk. Phos.](#), [DyLight®488](#), [DyLight®680](#),

[DyLight®800](#), [FITC](#), [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA1209\)](#)

North & South Tel: +1 800 265 7376

Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21

America Fax: +1 919 878 3751

Fax: +44 (0)1865 852 739

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M367616:200529'

Printed on 23 Mar 2021

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)