

## Datasheet: MCA349R

| Description:  | RAT ANTI HUMAN CD52 |
|---------------|---------------------|
| Specificity:  | CD52                |
| Other names:  | CAMPATH-1           |
| Format:       | Purified            |
| Product Type: | Monoclonal Antibody |
| Clone:        | YTH66.9HL           |
| Isotype:      | IgM                 |
| Quantity:     | 0.1 ml              |
|               |                     |

## **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

|                            | Yes | No | Not Determined | Suggested Dilution |
|----------------------------|-----|----|----------------|--------------------|
| Flow Cytometry             |     |    |                | 1/20               |
| Immunohistology - Frozen   | •   |    |                | 1/20               |
| Immunohistology - Paraffin |     |    | •              |                    |
| ELISA                      |     |    |                |                    |
| Immunoprecipitation        |     |    |                |                    |
| Western Blotting           |     |    |                |                    |
| Cytotoxic Assays           | •   |    |                |                    |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

| Target Species              | Human   |
|-----------------------------|---|
| Species Cross<br>Reactivity | Reacts with: Cynomolgus monkey  N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. |
| Product Form                | Ig fraction - liquid  |
| Preparation                 | Ig fraction prepared by ammonium sulphate precipitation from tissue culture supernatant.  |

| Buffer Solution                | Phosphate buffered saline  |  |  |
|--------------------------------|--|--|--|
| Preservative<br>Stabilisers    | <0.1% Sodium Azide (NaN <sub>3</sub> )   |  |  |
| Approx. Protein Concentrations | IgM concentration 2 mg/ml  |  |  |
| Immunogen                      | Human T lymphocytes.   |  |  |
| External Database              |  |  |  |
| Links                          | UniProt:   |  |  |
|                                | P31358 Related reagents  |  |  |
|                                | Entrez Gene:   |  |  |
|                                | 1043 CD52 Related reagents   |  |  |
| Synonyms                       | CDW52, HE5   |  |  |
| RRID                           | AB_324105  |  |  |
| Fusion Partners                | Spleen cells from an immunized DA rat were fused with cells of the Y3/Ag.1.2.3 myeloma cell line.  |  |  |
| Specificity                    | Rat anti Human CD52 antibody, clone YTH66.9HL recognizes human CD52, also known as CAMPATH-1. The CD52 antigen is a small heavily glycosylated peptide attached to the cell surface membrane via a GPI link. The apparent molecular mass of the antigen on SDS-PAGE is 25-29 kDa.  |  |  |
|                                | CD52 is expressed at high density by lymphocytes, monocytes, eosinophils, thymocytes and macrophages. It is expressed by most lymphoid derived malignancies, although expression on myeloma cells is variable.   |  |  |
|                                | Humanized versions of CAMPATH-1 specific antibodies are used for the treatment of a range of lymphoid malignancies (Blatt et al. 2014).  |  |  |
| Flow Cytometry                 | Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.  |  |  |
| References                     | Hale, G. <i>et al.</i> (1985) Reactivity of rat monoclonal antibody CAMPATH-1 with human leukaemia cells and its possible application for autologous bone marrow transplantation.  Br J Hale G. <i>et al.</i> (1998) Improving the outcome of bone marrow transplantation by using |  |  |
|                                | 2. Hale, G. <i>et al.</i> (1998) Improving the outcome of bone marrow transplantation by using CD52 monoclonal antibodies to prevent graft-versus-host disease and graft rejection.  Blood. 92 (12): 4581-90.  |  |  |
|                                | 3. Hale, G. <i>et al.</i> (1983) Effects of monoclonal anti-lymphocyte antibodies <i>in vivo</i> in  |  |  |
|                                | monkeys and humans. Mol Biol Med. 1 (3): 321-34.   |  |  |
|                                | 4. van der Windt, D.J. <i>et al.</i> (2010) Investigation of lymphocyte depletion and repopulation using alemtuzumab (Campath-1H) in cynomolgus monkeys. <u>Am J Transplant. 2010 Apr;10(4):773-83.</u>  |  |  |

|                                  | 5. van Wanrooij, R.L.J. <i>et al.</i> (2021) Adult-Onset Autoimmune I<br>Tertiary Referral Center. <u>Clin Transl Gastroenterol.</u> 12 (8): e003   |                      |
|----------------------------------|---|----------------------|
| Storage                          | This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. |                      |
|                                  | Avoid repeated freezing and thawing as this may denature the frost-free freezers is not recommended.  | antibody. Storage in |
| Guarantee                        | 12 months from date of despatch   |                      |
| Health And Safety<br>Information | Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA349R">https://www.bio-rad-antibodies.com/SDS/MCA349R</a> 10040  |                      |
| Regulatory                       | For research purposes only  |                      |

### **Related Products**

# **Recommended Secondary Antibodies**

Goat Anti Rat IgM (302001...) <u>Biotin</u> Goat Anti Rat IgM (STAR116...) <u>HRP</u>

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody\_sales\_us@bio-rad.com

 ${\bf Email: antibody\_sales\_uk@bio-rad.com}$ 

Email: antibody\_sales\_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M429871:240501'

#### Printed on 01 May 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint