

## Datasheet: MCA421

**BATCH NUMBER 149714**

<b>Description:</b>	RAT ANTI MOUSE IgG2a HEAVY CHAIN
<b>Specificity:</b>	IgG2a HEAVY CHAIN
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	LO-MG2a-7
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.25 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			5ug/ml as coating antibody
Immunoprecipitation			▪	
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Mouse
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.1% Sodium Azide
<b>Approx. Protein</b>	IgG concentration 1.0mg/ml

## Concentrations

---

Immunogen	Purified IgG from BALB/c mice.
-----------	--------------------------------

---

## External Database Links

### UniProt:

[P01863](#) [Related reagents](#)

[P01865](#) [Related reagents](#)

[P01864](#) [Related reagents](#)

### Entrez Gene:

[380793](#) Igh-1a [Related reagents](#)

[380793](#) Igh-1a [Related reagents](#)

[380793](#) Igh-1a [Related reagents](#)

---

RRID	AB_321822
------	-----------

---

Fusion Partners	Spleen cells from immunised LOU/c rats were fused with cells of the rat IR983F myeloma cell line.
-----------------	---

---

Specificity	<b>Rat anti Mouse IgG2a:Biotin, clone LO-MG2a-7</b> recognizes the gamma 2a heavy chain of mouse immunoglobulin and does not cross-react with other murine immunoglobulin classes or subclasses. Rat anti Mouse IgG2a Heavy Chain antibody, clone LO-MG2a-7 recognizes an allotypic determinant upon mouse IgG2a. It recognizes the IgH1a allotype (as expressed in Balb/c mice), but not the IgH1b allotype (as expressed in C57/BL mice).
-------------	---

---

## References

1. Jarman, E.R. & Lamb, J.R. (2004) Reversal of established CD4+ type 2 T helper-mediated allergic airway inflammation and eosinophilia by therapeutic treatment with DNA vaccines limits progression towards chronic inflammation and remodelling. [Immunology. 112 \(4\): 631-42.](#)
2. Bazin, H. (1982) Production of rat monoclonal antibodies with the LOU rat non secreting IR983F myeloma cell line. Prot Biol Fluids 615 - 8. Pergamon Press, Oxford and New York.
3. Nagao, K. *et al.* (2003) Role of prostaglandin I2 in airway remodeling induced by repeated allergen challenge in mice. [Am J Respir Cell Mol Biol. 29: 314-20.](#)
4. Bazin, H. *et al.* (1984) Rat monoclonal antibodies. I. Rapid purification from in vitro culture supernatants. [J Immunol Methods. 66 \(2\): 261-9.](#)
5. Bazin, H. *et al.* (1984) Rat monoclonal antibodies. II. A rapid and efficient method of purification from ascitic fluid or serum. [J Immunol Methods. 71 \(1\): 9-16.](#)
6. Ramos, J.D.A. *et al.* (2009) Characterization of Biot 11 Monoclonal Antibodies with Constant Region Mutations Phil Sci Lett 2: 38- 48.
7. Ormstad, H. *et al.* (2003) The effect of endotoxin on the production of IgE, IgG1 and IgG2a antibodies against the cat allergen Fel d 1 in mice [Toxicology. 188: 309-18.](#)
8. Ormstad, H. *et al.* (2000) The fungal cell wall component beta-1,3-glucan has an adjuvant effect on the allergic response to ovalbumin in mice. [J Toxicol Environ Health A. 61: 55-67.](#)
9. Hall, G. *et al.* (2003) Suppression of allergen reactive Th2 mediated responses and pulmonary eosinophilia by intranasal administration of an immunodominant peptide is

linked to IL-10 production. [Vaccine. 21: 549-61.](#)

10. Tan, L.K. *et al.* (2006) Intramuscular immunization with DNA construct containing Der p 2 and signal peptide sequences primed strong IgE production. [Vaccine. 24: 5762-71.](#)

11. Instanes, C. and Hetland, G. (2004) Deoxynivalenol (DON) is toxic to human colonic, lung and monocytic cell lines, but does not increase the IgE response in a mouse model for allergy. [Toxicology. 204: 13-21.](#)

12. Liedén, A. *et al.* (2009) Cornulin, a marker of late epidermal differentiation, is down-regulated in eczema. [Allergy. 64:304-11.](#)

13. Wolfowicz, C.B. *et al.* (2003) Expression and immunogenicity of the major house dust mite allergen Der p 1 following DNA immunization. [Vaccine. 21: 1195-204.](#)

14. Hayes, K.S. *et al.* (2017) Chronic *Trichuris muris* infection causes neoplastic change in the intestine and exacerbates tumour formation in APC min/+ mice. [PLoS Negl Trop Dis. 11 \(6\): e0005708.](#)

15. DeGiovanni, C. *et al.* (2019) Cancer Vaccines Co-Targeting HER2/Neu and IGF1R. [Cancers \(Basel\). 11 \(4\) Apr 11 \[Epub ahead of print\].](#)

---

<b>Further Reading</b>	1. Querinjean, P. <i>et al.</i> (1972) Transplantable immunoglobulin-secreting tumours in rats. Purification and chemical characterization of four kappa chains from LOU-Wsl rats. <a href="#">Eur J Biochem. 31 (2): 354-9.</a>
------------------------	--

---

<b>Storage</b>	Store at +4°C or at -20°C if preferred.
----------------	---

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

<b>Guarantee</b>	12 months from date of despatch
------------------	---------------------------------

---

<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA421">https://www.bio-rad-antibodies.com/SDS/MCA421</a> 10040
--------------------------------------	---

---

<b>Regulatory</b>	For research purposes only
-------------------	----------------------------

---

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	<a href="#">DyLight@800</a>
Rabbit Anti Rat IgG (STAR17...)	<a href="#">FITC</a>
Goat Anti Rat IgG (STAR69...)	<a href="#">FITC</a>
Goat Anti Rat IgG (STAR73...)	<a href="#">RPE</a>
Rabbit Anti Rat IgG (STAR21...)	<a href="#">HRP</a>
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	<a href="#">DyLight@550</a> , <a href="#">DyLight@650</a> , <a href="#">DyLight@800</a>
Goat Anti Rat IgG (STAR131...)	<a href="#">Alk. Phos.</a> , <a href="#">Biotin</a>
Goat Anti Rat IgG (STAR72...)	<a href="#">HRP</a>

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M367674:200529'

Printed on 31 Jan 2025

---

© 2025 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)