

Datasheet: MCA43FT

BATCH NUMBER 158831

Description:	MOUSE ANTI RAT CD45:FITC
Specificity:	CD45
Other names:	LCA
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	OX-1
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Rat		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide		
Stabilisers	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml		
Immunogen	Rat thymocyte membrane glycoproteins.		

External Database Links	UniProt: P04157 Related reagents Entrez Gene: 24699 Ptprc Related reagents
RRID	AB_322580
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the NS1 mouse myeloma cell line.
Specificity	<p>Mouse anti Rat CD45 antibody, clone OX-1 recognizes CD45, also known as the leucocyte common antigen (LCA). The leucocyte common antigen consists of a family of heavily glycosylated membrane glycoproteins of molecular weight 180 – 240kDa.</p> <p>Antibodies recognising a common epitope on all of these isoforms are termed CD45, whilst those recognising only individual isoforms are termed CD45RA, CD45RO etc. OX-1 reacts with all forms of CD45 expressed by all haematopoietic cells, except erythrocytes.</p> <p>CD45 isoforms play complex roles in T-cell and B-cell antigen receptor signal transduction.</p> <p>This product is routinely tested in flow cytometry on rat splenocytes</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> Sunderland, C.A. <i>et al.</i> (1979) Purification with monoclonal antibody of a predominant leukocyte-common antigen and glycoprotein from rat thymocytes. Eur J Immunol. 9 (2): 155-9. Woollett, G.R. <i>et al.</i> (1985) Molecular and antigenic heterogeneity of the rat leukocyte-common antigen from thymocytes and T and B lymphocytes. Eur J Immunol. 15 (2): 168-73. Martín, A. <i>et al.</i> (1995) Passive dual immunization against tumour necrosis factor-alpha (TNF-alpha) and IL-1 beta maximally ameliorates acute aminonucleoside nephrosis. Clin Exp Immunol. 99 (2): 283-8. Sato, K. <i>et al.</i> (2001) Carbon monoxide generated by heme oxygenase-1 suppresses the rejection of mouse-to-rat cardiac transplants. J Immunol. 166 (6): 4185-94. Murakami, K. <i>et al.</i> (2000) Regulation of mast cell signaling through high-affinity IgE receptor by CD45 protein tyrosine phosphatase. Int Immunol. 12 (2): 169-76. Standring, R. <i>et al.</i> (1978) The predominant heavily glycosylated glycoproteins at the surface of rat lymphoid cells are differentiation antigens. Eur J Immunol. 8 (12): 832-9. Giezeman-Smits, K.M. <i>et al.</i> (1999) The regulatory role of CD45 on rat NK cells in target cell lysis. J Immunol. 163 (1): 71-6. Zilka, N. <i>et al.</i> (2009) Human misfolded truncated tau protein promotes activation of microglia and leukocyte infiltration in the transgenic rat model of tauopathy. J Neuroimmunol. 209 (1-2): 16-25. Schupp, N. <i>et al.</i> (2011) Mineralocorticoid receptor-mediated DNA damage in kidneys of

- DOCA-salt hypertensive rats. [FASEB J. 25 \(3\): 968-78.](#)
10. Ermert, L. *et al.* (2001) Comparison of different detection methods in quantitative microdensitometry. [Am J Pathol. 158: 407-17.](#)
 11. Jeong, H.K. *et al.* (2010) Inflammatory responses are not sufficient to cause delayed neuronal death in ATP-induced acute brain injury. [PLoS One. 5: e13756.](#)
 12. Leonardo, C.C. *et al.* (2009) Inhibition of gelatinase activity reduces neural injury in an ex vivo model of hypoxia-ischemia. [Neuroscience. 160: 755-66.](#)
 13. Markusic, D.M. *et al.* (2010) Separating lentiviral vector injection and induction of gene expression in time, does not prevent an immune response to rTA in rats. [PLoS One. 5: e9974.](#)
 14. Vaschetto, R. *et al.* (2010) Renal hypoperfusion and impaired endothelium-dependent vasodilation in an animal model of VILI: the role of the peroxynitrite-PARP pathway [Crit Care. 14: R45.](#)
 15. Ladhoff, J. *et al.* (2010) Immune privilege of endothelial cells differentiated from endothelial progenitor cells. [Cardiovasc Res. 88: 121-9.](#)
 16. Yao, Y. *et al.* (2016) Alendronate Attenuates Spinal Microglial Activation and Neuropathic Pain. [J Pain. 17 \(8\): 889-903.](#)
 17. Wang, C. *et al.* (2015) Small activating RNA induces myogenic differentiation of rat adipose-derived stem cells by upregulating MyoD. [Int Braz J Urol. 41 \(4\): 764-72.](#)
 18. Tanner, D.C. *et al.* (2015) cFLIP is critical for oligodendrocyte protection from inflammation. [Cell Death Differ. 22 \(9\): 1489-501.](#)
 19. Runesson, E. *et al.* (2015) Nucleostemin- and Oct 3/4-positive stem/progenitor cells exhibit disparate anatomical and temporal expression during rat Achilles tendon healing. [BMC Musculoskelet Disord. 16: 212.](#)
 20. Hellenbrand, D.J. *et al.* (2019) Sustained interleukin-10 delivery reduces inflammation and improves motor function after spinal cord injury. [J Neuroinflammation. 16 \(1\): 93.](#)
 21. Pilipović, I. *et al.* (2019) Propranolol diminished severity of rat EAE by enhancing immunoregulatory/protective properties of spinal cord microglia. [Neurobiol Dis. Nov 2 \[Epub ahead of print\].](#)
 22. Collins, J.J.P. *et al.* (2018) Impaired Angiogenic Supportive Capacity and Altered Gene Expression Profile of Resident CD146⁺ Mesenchymal Stromal Cells Isolated from Hyperoxia-Injured Neonatal Rat Lungs. [Stem Cells Dev. 27 \(16\): 1109-24.](#)
 23. Porwal, K. *et al.* (2019) Increased bone marrow-specific adipogenesis by clofazimine causes impaired fracture healing, osteopenia and osteonecrosis without extra-skeletal effects in rats. [Toxicol Sci. kfz172.](#)
 24. Dabrowska, S. *et al.* (2021) Neuroinflammation evoked by brain injury in a rat model of lacunar infarct. [Exp Neurol. 336: 113531.](#)
 25. Elabi, O.F. *et al.* (2021) L-dopa-Dependent Effects of GLP-1R Agonists on the Survival of Dopaminergic Cells Transplanted into a Rat Model of Parkinson Disease. [Int J Mol Sci. 22\(22\):12346.](#)
 26. Hou, Y. *et al.* (2021) Pseudoginsenoside-F11 promotes functional recovery after transient cerebral ischemia by regulating the microglia/macrophage polarization in rats. [Int Immunopharmacol. 99: 107896.](#)
 27. Eweida, A. *et al.* (2022) Systemically injected bone marrow mononuclear cells specifically home to axially vascularized tissue engineering constructs. [PLoS One. 17 \(8\): e0272697.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA43FT>
10041

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA1209F\)](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M385846:210513'

Printed on 05 Feb 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)