

## Datasheet: MCA43GA

**BATCH NUMBER 1709**

<b>Description:</b>	MOUSE ANTI RAT CD45
<b>Specificity:</b>	CD45
<b>Other names:</b>	LCA
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	OX-1
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/100
Immunohistology - Frozen (1)	▪			
Immunohistology - Paraffin		▪		
ELISA			▪	
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

**(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.**

<b>Target Species</b>	Rat
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline

<b>Preservative Stabilisers</b>	0.09% Sodium Azide
<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Rat thymocyte membrane glycoproteins.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P04157</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">24699</a> Ptprc    <a href="#">Related reagents</a></p>
<b>RRID</b>	AB_566759
<b>Fusion Partners</b>	Spleen cells from immunised BALB/c mice were fused with cells of the NS1 mouse myeloma cell line.
<b>Specificity</b>	<p><b>Mouse anti Rat CD45 antibody, clone OX-1</b> recognizes CD45, also known as the leucocyte common antigen (LCA). The leucocyte common antigen consists of a family of heavily glycosylated membrane glycoproteins of molecular weight 180 – 240kDa.</p> <p>Antibodies recognising a common epitope on all of these isoforms are termed CD45, whilst those recognising only individual isoforms are termed CD45RA, CD45RO etc. OX-1 reacts with all forms of CD45 expressed by all haematopoietic cells, except erythrocytes.</p> <p>CD45 isoforms play complex roles in T-cell and B-cell antigen receptor signal transduction.</p> <p>This product is routinely tested in flow cytometry on rat splenocytes</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
<b>References</b>	<ol style="list-style-type: none"> <li>1. Sunderland, C.A. <i>et al.</i> (1979) Purification with monoclonal antibody of a predominant leukocyte-common antigen and glycoprotein from rat thymocytes. <a href="#">Eur J Immunol. 9 (2): 155-9.</a></li> <li>2. Woollett, G.R. <i>et al.</i> (1985) Molecular and antigenic heterogeneity of the rat leukocyte-common antigen from thymocytes and T and B lymphocytes. <a href="#">Eur J Immunol. 15 (2): 168-73.</a></li> <li>3. Martín, A. <i>et al.</i> (1995) Passive dual immunization against tumour necrosis factor-alpha (TNF-alpha) and IL-1 beta maximally ameliorates acute aminonucleoside nephrosis. <a href="#">Clin Exp Immunol. 99 (2): 283-8.</a></li> <li>4. Sato, K. <i>et al.</i> (2001) Carbon monoxide generated by heme oxygenase-1 suppresses the rejection of mouse-to-rat cardiac transplants. <a href="#">J Immunol. 166 (6): 4185-94.</a></li> <li>5. Murakami, K. <i>et al.</i> (2000) Regulation of mast cell signaling through high-affinity IgE</li> </ol>

- receptor by CD45 protein tyrosine phosphatase. [Int Immunol. 12 \(2\): 169-76.](#)
6. Standring, R. *et al.* (1978) The predominant heavily glycosylated glycoproteins at the surface of rat lymphoid cells are differentiation antigens. [Eur J Immunol. 8 \(12\): 832-9.](#)
7. Giezeman-Smits, K.M. *et al.* (1999) The regulatory role of CD45 on rat NK cells in target cell lysis. [J Immunol. 163 \(1\): 71-6.](#)
8. Zilka, N. *et al.* (2009) Human misfolded truncated tau protein promotes activation of microglia and leukocyte infiltration in the transgenic rat model of tauopathy. [J Neuroimmunol. 209 \(1-2\): 16-25.](#)
9. Schupp, N. *et al.* (2011) Mineralocorticoid receptor-mediated DNA damage in kidneys of DOCA-salt hypertensive rats. [FASEB J. 25 \(3\): 968-78.](#)
10. Ermert, L. *et al.* (2001) Comparison of different detection methods in quantitative microdensitometry. [Am J Pathol. 158: 407-17.](#)
11. Jeong, H.K. *et al.* (2010) Inflammatory responses are not sufficient to cause delayed neuronal death in ATP-induced acute brain injury. [PLoS One. 5: e13756.](#)
12. Leonardo, C.C. *et al.* (2009) Inhibition of gelatinase activity reduces neural injury in an ex vivo model of hypoxia-ischemia. [Neuroscience. 160: 755-66.](#)
13. Markusic, D.M. *et al.* (2010) Separating lentiviral vector injection and induction of gene expression in time, does not prevent an immune response to rtTA in rats. [PLoS One. 5: e9974.](#)
14. Vaschetto, R. *et al.* (2010) Renal hypoperfusion and impaired endothelium-dependent vasodilation in an animal model of VILI: the role of the peroxynitrite-PARP pathway [Crit Care. 14: R45.](#)
15. Ladhoff, J. *et al.* (2010) Immune privilege of endothelial cells differentiated from endothelial progenitor cells. [Cardiovasc Res. 88: 121-9.](#)
16. Yao, Y. *et al.* (2016) Alendronate Attenuates Spinal Microglial Activation and Neuropathic Pain. [J Pain. 17 \(8\): 889-903.](#)
17. Wang, C. *et al.* (2015) Small activating RNA induces myogenic differentiation of rat adipose-derived stem cells by upregulating MyoD. [Int Braz J Urol. 41 \(4\): 764-72.](#)
18. Tanner, D.C. *et al.* (2015) cFLIP is critical for oligodendrocyte protection from inflammation. [Cell Death Differ. 22 \(9\): 1489-501.](#)
19. Runesson, E. *et al.* (2015) Nucleostemin- and Oct 3/4-positive stem/progenitor cells exhibit disparate anatomical and temporal expression during rat Achilles tendon healing. [BMC Musculoskelet Disord. 16: 212.](#)
20. Hellenbrand, D.J. *et al.* (2019) Sustained interleukin-10 delivery reduces inflammation and improves motor function after spinal cord injury. [J Neuroinflammation. 16 \(1\): 93.](#)
21. Pilipović, I. *et al.* (2019) Propranolol diminished severity of rat EAE by enhancing immunoregulatory/protective properties of spinal cord microglia. [Neurobiol Dis. Nov 2 \[Epub ahead of print\].](#)

---

**Storage**

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

<b>Guarantee</b>	12 months from date of despatch
<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA43GA">https://www.bio-rad-antibodies.com/SDS/MCA43GA</a> 10040
<b>Regulatory</b>	For research purposes only

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	<a href="#">RPE</a>
Goat Anti Mouse IgG IgA IgM (STAR87...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (STAR76...)	<a href="#">RPE</a>
Goat Anti Mouse IgG (STAR70...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (H/L) (STAR117...)	<a href="#">Alk. Phos.</a> , <a href="#">DyLight@488</a> , <a href="#">DyLight@550</a> , <a href="#">DyLight@650</a> , <a href="#">DyLight@680</a> , <a href="#">DyLight@800</a> , <a href="#">FITC</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR13...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (Fc) (STAR120...)	<a href="#">FITC</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR9...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (STAR77...)	<a href="#">HRP</a>

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA1209\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)  
'M367696:200529'

Printed on 03 Sep 2024