

Datasheet: MCA453F

Description:	MOUSE ANTI RAT TCR ALPHA/BETA:FITC
Specificity:	TCR ALPHA/BETA
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	R73
Isotype:	lgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			Neat - 1/5

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Rat					
Species Cross Reactivity	Reacts with: Monkey, Cynomolgus monkey N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications of personal communications from the originators. Please refer to references indicated further information.					
Product Form	Purified IgG conju	ugated to Fluorescein Isotl	hiocyanate Isomer 1 (FITC) - liquid			
Max Ex/Em	Fluorophore FITC	Excitation Max (nm) 490	Emission Max (nm) 525			
Preparation	Purified IgG prepared supernatant	ared by affinity chromatog	raphy on Protein A from tissue culture			
Buffer Solution	Phosphate buffer	ed saline				

Preservative	0.09% Sodium Azide
Stabilisers	1% Bovine Serum Albumin
	170 Bovine Gerun Albumin
Approx. Protein	IgG concentration 0.1 mg/ml
Concentrations	igo concentration o. i mg/mi
Immunogen	Rat T blasts and erythrocytes.
RRID	AB_322439
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the X63.Ag8.653 mouse myeloma cell line.
Specificity	Mouse anti Rat TCR alpha/beta antibody, clone R73 recognizes a constant determinant on the beta chain of the rat alpha/beta T cell receptor, expressed by 97% of peripheral rat T cells as defined by the OX-52 marker. R73 is mitogenic for unseparated spleen cells and for purified T cells. In the rat thymus, mature medullary cells express the R73 determinant at the same levels as peripheral T cells, whereas 94% of CD4 - CD8 - thymocytes are R73 negative.
	Mouse anti Rat TCR alpha/beta antibody, clone R73 is reported to stimulate adhesion between Thymic Dendritic Cells and Thymocytes (Colic et al. 2010).
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ lymphocytes in 100ul.
References	1. Tomida, S. <i>et al.</i> (1994) Intercellular adhesion molecule-1 and leukocyte function-
	associated antigen-1 are involved in protection mediated by CD3+TCR alpha beta- T cells at the early stage after infection with Listeria monocytogenes in rats. Int Immunol. 6 (7):
	955-61.
	2. Colić, M. <i>et al.</i> (1996) Mechanisms involved in the binding of thymocytes to rat thymic
	dendritic cells. Dev Immunol. 5 (1): 37-51.
	3. Kanellis, J. et al. (2010) JNK signalling in human and experimental renal
	ischaemia/reperfusion injury. Nephrol Dial Transplant. 25: 2898-908.
	4. Nave, H. et al. (2008) Resistance of Janus kinase-2 dependent leptin signaling in
	natural killer (NK) cells: a novel mechanism of NK cell dysfunction in diet-induced obesity.
	Endocrinology. 149: 3370-8.
	5. Tsuchida, M. et al. (1994) Identification of CD4- CD8- alpha beta T cells in the
	subgrachnoid space of rats with experimental autoimmune encephalomyelitis. A possible

- 5. Tsuchida, M. *et al.* (1994) Identification of CD4- CD8- alpha beta T cells in the subarachnoid space of rats with experimental autoimmune encephalomyelitis. A possible route by which effector cells invade the lesions. lmmunology.81 (3): 420-7.
- 6. Matsumoto, Y. *et al.* (1994) Successful prevention and treatment of autoimmune encephalomyelitis by short-term administration of anti-T-cell receptor alpha beta antibody. Immunology. 81 (1): 1-7.
- 7. Pilipović, I. *et al.* (2010) Glucocorticoids, master modulators of the thymic catecholaminergic system? <u>Braz J Med Biol Res.</u> 43 (3): 279-84.
- 8. Milicevic, N.M. *et al.* (2005) T cells are required for the peripheral phase of B-cell maturation. <u>Immunology</u>. 116: 308-17.
- 9. Kenny, E. *et al.* (2000) Phenotypic analysis of peripheral CD4+ CD8+ T cells in the rat. <u>Immunology. 101: 178-84.</u>

- 10. Petrovic-Dergovic, D.M. *et al.* (2004) Somatostatin-14 alters the thymus size and relation among the thymocyte subpopulations in peripubertal rats. <u>Neuropeptides. 38:</u> 25-34.
- 11. Trinh, L. *et al.* (2008) The corneal endothelium in an endotoxin-induced uveitis model: correlation between in vivo confocal microscopy and immunohistochemistry. <u>Mol Vis. 14:</u> 1149-56.
- 12. Khalife, S. *et al.* (2016) Relationship Between *Pneumocystis carinii* Burden and the Degree of Host Immunosuppression in an Airborne Transmission Experimental Model. <u>J Eukaryot Microbiol.</u> 63 (3): 309-17.
- 13. Bat, E. *et al.* (2013) Physical properties and erosion behavior of poly(trimethylene carbonate-co-ε-caprolactone) networks. Macromol Biosci. 13 (5): 573-83.
- 14. Jörns, A. *et al.* (2015) TNF-α Antibody Therapy in Combination With the T-Cell-Specific Antibody Anti-TCR Reverses the Diabetic Metabolic State in the LEW.1AR1-iddm Rat. <u>Diabetes. 64 (8): 2880-91.</u>
- 15. Ahn, M. *et al.* (2015) Immunohistochemical study of Krüppel-like factor 4 in the spinal cords of rats with experimental autoimmune encephalomyelitis. <u>Acta Histochem. 117 (6):</u> 521-7.
- 16. Jörns A *et al.* (2014) Anti-TCR therapy combined with fingolimod for reversal of diabetic hyperglycemia by β cell regeneration in the LEW.1AR1-iddm rat model of type 1 diabetes. J Mol Med (Berl). 92 (7): 743-55.
- 17. Amos, L.A. *et al.* (2018) ASK1 inhibitor treatment suppresses p38/JNK signalling with reduced kidney inflammation and fibrosis in rat crescentic glomerulonephritis. <u>J Cell Mol Med.</u> 22 (9): 4522-33.
- 18. Koppe, C. *et al.* (2021) Local Inflammatory Response after Intramuscularly Implantation of Anti-Adhesive Plasma-Fluorocarbon-Polymer Coated Ti6Al4V Discs in Rats. Polymers (Basel). 13 (16): 2684.
- 19. Schmiedl, A. *et al.* (2021) Lung development and immune status under chronic LPS exposure in rat pups with and without CD26/DPP4 deficiency. <u>Cell Tissue Res. Oct 04</u> [Epub ahead of print].
- 20. Köhler, R. *et al.* (2022) Association of systemic antibody response against polyethylene terephthalate with inflammatory serum cytokine profile following implantation of differently coated vascular prostheses in a rat animal model. <u>J Biomed Mater Res A.</u> 110 (1): 52-63.
- 21. Martin, A. *et al.* (2018) Tumor-derived granzyme B-expressing neutrophils acquire antitumor potential after lipid A treatment. <u>Oncotarget. 9 (47): 28364-78.</u>
- 22. Onaru, K. *et al.* (2020) Immunotoxicity evaluation by subchronic oral administration of clothianidin in Sprague-Dawley rats. <u>J Vet Med Sci. 82 (3): 360-72.</u>
- 23. Midavaine, É. *et al.* (2024) Discovery of a CCR2-targeting pepducin therapy for chronic pain. <u>Pharmacol Res.</u>: 107242.
- 24. Jörns, A. *et al.* (2020) Translation of curative therapy concepts with T cell and cytokine antibody combinations for type 1 diabetes reversal in the IDDM rat. <u>J Mol Med (Berl). 98</u> (8): 1125-37.

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA453F 10041
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:FITC (MCA1209F)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M421881:230803'

Printed on 15 Apr 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint