

## Datasheet: MCA453G

<b>Description:</b>	MOUSE ANTI RAT TCR ALPHA/BETA
<b>Specificity:</b>	TCR ALPHA/BETA
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	R73
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.25 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen	▪			1/10 - 1/100
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Rat
<b>Species Cross Reactivity</b>	<p>Reacts with: Monkey, Cynomolgus monkey</p> <p><b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline

<b>Preservative Stabilisers</b>	0.09% Sodium Azide
<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Rat T Blasts and Erythrocytes.
<b>RRID</b>	AB_322435
<b>Fusion Partners</b>	Spleen cells from immunised BALB/c mice were fused with cells of the X63.Ag8.653 mouse myeloma cell line.
<b>Specificity</b>	<p><b>Mouse anti Rat TCR alpha/beta antibody, clone R73</b> recognizes a constant determinant on the beta chain of the rat alpha/beta T cell receptor, expressed by 97% of peripheral rat T cells as defined by the OX-52 marker. R73 is mitogenic for unseparated spleen cells and for purified T cells. In the rat thymus, mature medullary cells express the R73 determinant at the same levels as peripheral T cells, whereas 94% of CD4 - CD8 - thymocytes are R73 negative.</p> <p>Mouse anti Rat TCR alpha/beta antibody, clone R73 is reported to stimulate adhesion between Thymic Dendritic Cells and Thymocytes (<a href="#">Colic et al. 2010</a>).</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 100ul of whole blood or 10 <sup>6</sup> cells in 100ul.
<b>References</b>	<ol style="list-style-type: none"> <li>Tomida, S. <i>et al.</i> (1994) Intercellular adhesion molecule-1 and leukocyte function-associated antigen-1 are involved in protection mediated by CD3+TCR alpha beta- T cells at the early stage after infection with <i>Listeria monocytogenes</i> in rats. <a href="#">Int Immunol. 6 (7): 955-61.</a></li> <li>Colić, M. <i>et al.</i> (1996) Mechanisms involved in the binding of thymocytes to rat thymic dendritic cells. <a href="#">Dev Immunol. 5 (1): 37-51.</a></li> <li>Kanellis, J. <i>et al.</i> (2010) JNK signalling in human and experimental renal ischaemia/reperfusion injury. <a href="#">Nephrol Dial Transplant. 25: 2898-908.</a></li> <li>Nave, H. <i>et al.</i> (2008) Resistance of Janus kinase-2 dependent leptin signaling in natural killer (NK) cells: a novel mechanism of NK cell dysfunction in diet-induced obesity. <a href="#">Endocrinology. 149: 3370-8.</a></li> <li>Tsuchida, M. <i>et al.</i> (1994) Identification of CD4- CD8- alpha beta T cells in the subarachnoid space of rats with experimental autoimmune encephalomyelitis. A possible route by which effector cells invade the lesions. <a href="#">Immunology. 81 (3): 420-7.</a></li> <li>Matsumoto, Y. <i>et al.</i> (1994) Successful prevention and treatment of autoimmune encephalomyelitis by short-term administration of anti-T-cell receptor alpha beta antibody. <a href="#">Immunology. 81 (1): 1-7.</a></li> <li>Pilipović, I. <i>et al.</i> (2010) Glucocorticoids, master modulators of the thymic catecholaminergic system? <a href="#">Braz J Med Biol Res. 43 (3): 279-84.</a></li> <li>Milicevic, N.M. <i>et al.</i> (2005) T cells are required for the peripheral phase of B-cell</li> </ol>

- maturation. [Immunology. 116: 308-17.](#)
9. Kenny, E. *et al.* (2000) Phenotypic analysis of peripheral CD4+ CD8+ T cells in the rat. [Immunology. 101: 178-84.](#)
  10. Petrovic-Dergovic, D.M. *et al.* (2004) Somatostatin-14 alters the thymus size and relation among the thymocyte subpopulations in peripubertal rats. [Neuropeptides. 38: 25-34.](#)
  11. Trinh, L. *et al.* (2008) The corneal endothelium in an endotoxin-induced uveitis model: correlation between in vivo confocal microscopy and immunohistochemistry. [Mol Vis. 14: 1149-56.](#)
  12. Khalife, S. *et al.* (2016) Relationship Between *Pneumocystis carinii* Burden and the Degree of Host Immunosuppression in an Airborne Transmission Experimental Model. [J Eukaryot Microbiol. 63 \(3\): 309-17.](#)
  13. Bat, E. *et al.* (2013) Physical properties and erosion behavior of poly(trimethylene carbonate-co-ε-caprolactone) networks. [Macromol Biosci. 13 \(5\): 573-83.](#)
  14. Jörns, A. *et al.* (2015) TNF-α Antibody Therapy in Combination With the T-Cell-Specific Antibody Anti-TCR Reverses the Diabetic Metabolic State in the LEW.1AR1-iddm Rat. [Diabetes. 64 \(8\): 2880-91.](#)
  15. Ahn, M. *et al.* (2015) Immunohistochemical study of Krüppel-like factor 4 in the spinal cords of rats with experimental autoimmune encephalomyelitis. [Acta Histochem. 117 \(6\): 521-7.](#)
  16. Jörns A *et al.* (2014) Anti-TCR therapy combined with fingolimod for reversal of diabetic hyperglycemia by β cell regeneration in the LEW.1AR1-iddm rat model of type 1 diabetes. [J Mol Med \(Berl\). 92 \(7\): 743-55.](#)
  17. Amos, L.A. *et al.* (2018) ASK1 inhibitor treatment suppresses p38/JNK signalling with reduced kidney inflammation and fibrosis in rat crescentic glomerulonephritis. [J Cell Mol Med. 22 \(9\): 4522-33.](#)
  18. Koppe, C. *et al.* (2021) Local Inflammatory Response after Intramuscularly Implantation of Anti-Adhesive Plasma-Fluorocarbon-Polymer Coated Ti6Al4V Discs in Rats. [Polymers \(Basel\). 13 \(16\): 2684.](#)
  19. Schmiedl, A. *et al.* (2021) Lung development and immune status under chronic LPS exposure in rat pups with and without CD26/DPP4 deficiency. [Cell Tissue Res. Oct 04 \[Epub ahead of print\].](#)
  20. Köhler, R. *et al.* (2022) Association of systemic antibody response against polyethylene terephthalate with inflammatory serum cytokine profile following implantation of differently coated vascular prostheses in a rat animal model. [J Biomed Mater Res A. 110 \(1\): 52-63.](#)

<b>Storage</b>	<p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.</p>
<b>Guarantee</b>	12 months from date of despatch
<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA453G">https://www.bio-rad-antibodies.com/SDS/MCA453G</a>

**Regulatory**

For research purposes only

**Related Products****Recommended Secondary Antibodies**

Goat Anti Mouse IgG (STAR77...)	<a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR12...)	<a href="#">RPE</a>
Goat Anti Mouse IgG (STAR70...)	<a href="#">FITC</a>
Goat Anti Mouse IgG IgA IgM (STAR87...)	<a href="#">Alk. Phos.</a> , <a href="#">HRP</a>
Goat Anti Mouse IgG (STAR76...)	<a href="#">RPE</a>
Goat Anti Mouse IgG (H/L) (STAR117...)	<a href="#">Alk. Phos.</a> , <a href="#">DyLight®488</a> , <a href="#">DyLight®550</a> , <a href="#">DyLight®650</a> , <a href="#">DyLight®680</a> , <a href="#">DyLight®800</a> , <a href="#">FITC</a> , <a href="#">HRP</a>
Goat Anti Mouse IgG (Fc) (STAR120...)	<a href="#">FITC</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR9...)	<a href="#">FITC</a>
Rabbit Anti Mouse IgG (STAR13...)	<a href="#">HRP</a>

**Recommended Negative Controls**[MOUSE IgG1 NEGATIVE CONTROL \(MCA1209\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)  
'M384049:210513'

**Printed on 12 Aug 2023**