

Datasheet: MCA453GA

BATCH NUMBER 164445

Description:	MOUSE ANTI RAT TCR ALPHA/BETA
Specificity:	TCR ALPHA/BETA
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	R73
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen	▪			1/10 - 1/100
Immunohistology - Paraffin			▪	
Immunohistology - Resin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species

Rat

Species Cross Reactivity

Reacts with: Monkey, Cynomolgus monkey

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG - liquid

Preparation

Purified IgG prepared by affinity chromatography on Protein A from tissue culture

supernatant

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Rat T blasts and erythrocytes.
RRID	AB_567214
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the X63.Ag8.653 mouse myeloma cell line.
Specificity	<p>Mouse anti Rat TCR alpha/beta antibody, clone R73 recognizes a constant determinant on the beta chain of the rat alpha/beta T cell receptor, expressed by 97% of peripheral rat T cells as defined by the OX-52 marker. R73 is mitogenic for unseparated spleen cells and for purified T cells. In the rat thymus, mature medullary cells express the R73 determinant at the same levels as peripheral T cells, whereas 94% of CD4 - CD8 - thymocytes are R73 negative.</p> <p>Mouse anti Rat TCR alpha/beta antibody, clone R73 is reported to stimulate adhesion between Thymic Dendritic Cells and Thymocytes (Colic et al. 2010).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 100ul of whole blood or 10 ⁶ cells in 100ul. Method sheets are available on request.
References	<ol style="list-style-type: none">1. Tomida, S. <i>et al.</i> (1994) Intercellular adhesion molecule-1 and leukocyte function-associated antigen-1 are involved in protection mediated by CD3+TCR alpha beta- T cells at the early stage after infection with <i>Listeria monocytogenes</i> in rats. Int Immunol. 6 (7): 955-61.2. Colić, M. <i>et al.</i> (1996) Mechanisms involved in the binding of thymocytes to rat thymic dendritic cells. Dev Immunol. 5 (1): 37-51.3. Kanellis, J. <i>et al.</i> (2010) JNK signalling in human and experimental renal ischaemia/reperfusion injury. Nephrol Dial Transplant. 25: 2898-908.4. Nave, H. <i>et al.</i> (2008) Resistance of Janus kinase-2 dependent leptin signaling in natural killer (NK) cells: a novel mechanism of NK cell dysfunction in diet-induced obesity. Endocrinology. 149: 3370-8.5. Tsuchida, M. <i>et al.</i> (1994) Identification of CD4- CD8- alpha beta T cells in the subarachnoid space of rats with experimental autoimmune encephalomyelitis. A possible route by which effector cells invade the lesions. Immunology. 81 (3): 420-7.6. Matsumoto, Y. <i>et al.</i> (1994) Successful prevention and treatment of autoimmune encephalomyelitis by short-term administration of anti-T-cell receptor alpha beta antibody. Immunology. 81 (1): 1-7.

7. Pilipović, I. *et al.* (2010) Glucocorticoids, master modulators of the thymic catecholaminergic system? [Braz J Med Biol Res. 43 \(3\): 279-84.](#)
8. Milicevic, N.M. *et al.* (2005) T cells are required for the peripheral phase of B-cell maturation. [Immunology. 116: 308-17.](#)
9. Kenny, E. *et al.* (2000) Phenotypic analysis of peripheral CD4+ CD8+ T cells in the rat. [Immunology. 101: 178-84.](#)
10. Petrovic-Dergovic, D.M. *et al.* (2004) Somatostatin-14 alters the thymus size and relation among the thymocyte subpopulations in peripubertal rats. [Neuropeptides. 38: 25-34.](#)
11. Trinh, L. *et al.* (2008) The corneal endothelium in an endotoxin-induced uveitis model: correlation between in vivo confocal microscopy and immunohistochemistry. [Mol Vis. 14: 1149-56.](#)
12. Khalife, S. *et al.* (2016) Relationship Between *Pneumocystis carinii* Burden and the Degree of Host Immunosuppression in an Airborne Transmission Experimental Model. [J Eukaryot Microbiol. 63 \(3\): 309-17.](#)
13. Bat, E. *et al.* (2013) Physical properties and erosion behavior of poly(trimethylene carbonate-co- ϵ -caprolactone) networks. [Macromol Biosci. 13 \(5\): 573-83.](#)
14. Jörns, A. *et al.* (2015) TNF- α Antibody Therapy in Combination With the T-Cell-Specific Antibody Anti-TCR Reverses the Diabetic Metabolic State in the LEW.1AR1-iddm Rat. [Diabetes. 64 \(8\): 2880-91.](#)
15. Ahn, M. *et al.* (2015) Immunohistochemical study of Krüppel-like factor 4 in the spinal cords of rats with experimental autoimmune encephalomyelitis. [Acta Histochem. 117 \(6\): 521-7.](#)
16. Jörns A *et al.* (2014) Anti-TCR therapy combined with fingolimod for reversal of diabetic hyperglycemia by β cell regeneration in the LEW.1AR1-iddm rat model of type 1 diabetes. [J Mol Med \(Berl\). 92 \(7\): 743-55.](#)
17. Amos, L.A. *et al.* (2018) ASK1 inhibitor treatment suppresses p38/JNK signalling with reduced kidney inflammation and fibrosis in rat crescentic glomerulonephritis. [J Cell Mol Med. 22 \(9\): 4522-33.](#)
18. Koppe, C. *et al.* (2021) Local Inflammatory Response after Intramuscularly Implantation of Anti-Adhesive Plasma-Fluorocarbon-Polymer Coated Ti6Al4V Discs in Rats. [Polymers \(Basel\). 13 \(16\): 2684.](#)
19. Schmiedl, A. *et al.* (2021) Lung development and immune status under chronic LPS exposure in rat pups with and without CD26/DPP4 deficiency. [Cell Tissue Res. Oct 04 \[Epub ahead of print\].](#)
20. Köhler, R. *et al.* (2022) Association of systemic antibody response against polyethylene terephthalate with inflammatory serum cytokine profile following implantation of differently coated vascular prostheses in a rat animal model. [J Biomed Mater Res A. 110 \(1\): 52-63.](#)
21. Martin, A. *et al.* (2018) Tumor-derived granzyme B-expressing neutrophils acquire antitumor potential after lipid A treatment. [Oncotarget. 9 \(47\): 28364-78.](#)
22. Onaru, K. *et al.* (2020) Immunotoxicity evaluation by subchronic oral administration of clothianidin in Sprague-Dawley rats. [J Vet Med Sci. 82 \(3\): 360-72.](#)
23. Midavaine, É. *et al.* (2024) Discovery of a CCR2-targeting pepducin therapy for chronic pain. [Pharmacol Res. : 107242.](#)
24. Jörns, A. *et al.* (2020) Translation of curative therapy concepts with T cell and cytokine antibody combinations for type 1 diabetes reversal in the IDDM rat. [J Mol Med \(Berl\). 98](#)

[\(8\): 1125-37.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA453GA>
10040

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
Goat Anti Mouse IgG IgA IgM (STAR87...) [HRP](#)
Goat Anti Mouse IgG (STAR76...) [RPE](#)
Goat Anti Mouse IgG (STAR70...) [FITC](#)
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),
[FITC](#), [HRP](#)
Goat Anti Mouse IgG (STAR77...) [HRP](#)
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA1209\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M421882:230803'

Printed on 14 Jun 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)