

Datasheet: MCA45R

BATCH NUMBER 161299

Description:	MOUSE ANTI RAT MHC CLASS II RT1Bu/I
Specificity:	MHC CLASS II RT1Bu/I
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	OX-3
Isotype:	IgG1
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin	▪			
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Rat
Species Cross Reactivity	<p>Reacts with: Mouse</p> <p>N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Rat thymocyte membrane glycoproteins.
RRID	AB_322406
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells from the NS1 mouse myeloma cell line.
Specificity	<p>Mouse anti Rat MHC Class II RT1Bu/L antibody, clone OX-3 recognizes a polymorphic determinant of the rat RT1B MHC class II antigen, reacting with haplotypes u and l. The literature reports reactivity with Lewis, Wistar and AO strain rats but not BN, DA or PVG/c strains. This antibody is useful for distinguishing RT1B positive cells from different rat strains, e.g. for recognising cells of donor origin in bone marrow reconstituted radiation chimaeras.</p> <p>The major histocompatibility complex (MHC) is a cluster of genes that are important in the immune response to infections. In rats, this complex is referred to as the RT1 region. In mice, this complex is referred to as the H-2 region.</p> <p>Mouse anti Rat MHC Class II RT1Bu/L antibody, clone OX-3 also cross reacts with mouse strains of the H-2 haplotypes b and s. Analysis of recombinant mouse strains has mapped the OX-3 determinant to the H-2I-A region.</p> <p>This product is routinely tested in flow cytometry on Lewis rat splenocytes.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul
Immunohistology	<p>This product does not require protein digestion pre-treatment of paraffin embedded sections e.g. trypsin or pronase prior to staining.</p> <p>This product does not require antigen retrieval using heat treatment methods prior to staining of paraffin sections.</p>
References	<ol style="list-style-type: none"> 1. McMaster, W.R. & Williams, A.F. (1979) Identification of Ia glycoproteins in rat thymus and purification from rat spleen. Eur J Immunol. 9 (6): 426-33. 2. McMaster, W.R. & Williams, A.F. (1979) Monoclonal antibodies to Ia antigens from rat thymus: cross reactions with mouse and human and use in purification of rat Ia glycoproteins. Immunol Rev. 47: 117-37. 3. Barclay, A.N. & Mayrhofer, G. (1981) Bone marrow origin of Ia-positive cells in the medulla rat thymus. J Exp Med. 153 (6): 1666-71.

4. Zhang, J. *et al.* (1997) Expression of major histocompatibility complex molecules in rodent retina. Immunohistochemical study. [Invest Ophthalmol Vis Sci. 38 \(9\): 1848-57.](#)
5. Hahm, K.B. *et al.* (2000) Loss of TGF-beta signaling contributes to autoimmune pancreatitis. [J Clin Invest. 105 \(8\): 1057-65.](#)
6. Wu, S.Y. *et al.* (2016) Estrogen ameliorates microglial activation by inhibiting the Kir2.1 inward-rectifier K(+) channel. [Sci Rep. 6: 22864.](#)
7. Fisher, R.A. *et al.* (1996) Induction of long-term graft tolerance and donor/recipient chimerism. [J Surg Res. 60 \(1\): 181-5.](#)
8. Keller, R. *et al.* (1988) Modulation of major histocompatibility complex (MHC) expression by interferons and microbial agents. Independent regulation of MHC class II expression and induction of tumoricidal activity in bone marrow-derived mononuclear phagocytes. [Scand J Immunol. 28 \(1\): 113-21.](#)
9. Streit, W.J. *et al.* (1989) Peripheral nerve lesion produces increased levels of major histocompatibility complex antigens in the central nervous system. [J Neuroimmunol. 21 \(2-3\): 117-23.](#)
10. Roggin, K.K. *et al.* (2001) Macrophage phenotype during cholestatic injury and repair: the persistent inflammatory response. [J Pediatr Surg. 36 \(1\): 220-8.](#)
11. Reutzel-Selke A *et al.* (2003) Short-term immunosuppressive treatment of the donor ameliorates consequences of ischemia/ reperfusion injury and long-term graft function in renal allografts from older donors. [Transplantation. 75 \(11\): 1786-92.](#)
12. Heidenhain, C. *et al.* (2003) The impact of immune-activating processes following transplantation on chronic allograft nephropathy. [Kidney Int. 64 \(3\): 1125-33.](#)
13. Hahm, K.B. *et al.* (2001) Loss of transforming growth factor beta signalling in the intestine contributes to tissue injury in inflammatory bowel disease. [Gut. 49 \(2\): 190-8.](#)
14. Pascher A *et al.* (2006) Rat cytomegalovirus infection interferes with anti-CD4 mAb-(RIB 5/2) mediated tolerance and induces chronic allograft damage. [Am J Transplant. 6 \(9\): 2035-45.](#)
15. Hartmann CB *et al.* (2005) Immunotoxicity of gallium arsenide on antigen presentation: comparative study of intratracheal and intraperitoneal exposure routes. [J Immunotoxicol. 2 \(1\): 1-9.](#)
16. Lima, R.R. *et al.* (2022) Inflammatory Response and Secondary White Matter Damage to the Corpus Callosum after Focal Striatal Stroke in Rats. [Int J Mol Sci. 23 \(6\)Mar 16 \[Epub ahead of print\].](#)
17. Duhalde Vega, M *et al.* (2022) PD-1/PD-L1 blockade abrogates a dysfunctional innate-adaptive immune axis in critical β -coronavirus disease. [Sci Adv. 8 \(38\): eabn6545.](#)

Further Reading

1. Barclay, A.N. (1981) The localization of populations of lymphocytes defined by monoclonal antibodies in rat lymphoid tissues. [Immunology. 42 \(4\): 593-600.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee

12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at:
<https://www.bio-rad-antibodies.com/SDS/MCA45R>
10040

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...) [HRP](#)
Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
Goat Anti Mouse IgG (STAR70...) [FITC](#)
Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)
Goat Anti Mouse IgG (STAR76...) [RPE](#)
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),
[FITC](#), [HRP](#)
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA1209\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------	------------------	-------------------------------------------------------------------------------------------------------------------------------------------------	---------------	-----------------------------------------------------------------------------------------------------------------------------------------------------

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M384030:210513'

Printed on 07 Apr 2025

© 2025 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)