

Datasheet: MCA4610T BATCH NUMBER 154626

Description:	MOUSE ANTI HUMAN CD15	
Specificity:	CD15	
Other names:	LEWIS X	
Format:	Purified	
Product Type:	Monoclonal Antibody	
Clone:	HI98	
Isotype:	IgM	
Quantity:	25 μg	

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				1/5 - 1/10
Immunohistology - Frozen				
Immunohistology - Paraffin				
ELISA				
Immunoprecipitation				
Western Blotting				

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human	
Species Cross Reactivity	Reacts with: Chimpanzee N.B. Antibody reactivity and working conditions may vary betwee reactivity is derived from testing within our laboratories, peer-revie personal communications from the originators. Please refer to refurther information.	ewed publications or
Product Form	Purified IgM - liquid	

Preparation	Ig fraction prepared by affinity chromatography
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	Ig concentration 0.5mg/ml
RRID	AB_2247102
Specificity	Mouse anti Human CD15 antibody, clone HI98 recognises human CD15, otherwise known as Lewis X and stage-specfic embryonic antigen-1 (SSEA-1). The CD15 antigen itself a terminal trisaccharide, Gal-beta1-4 (Fuc-alpha1-3) GlcNAc (3-FAL), which is found on a variety of glycoproteins and glycolipids. CD15 is predominately expressed by peripheral blood granulocytes and various epithelial tissues. It plays an important role in cell adhesion and has been used as a marker for diagnosis of Hodgkin's lymphoma (Zukerberg et al. 1991).
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.
Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this madenature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA4610T 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgM (STAR138...) Alk. Phos.
Goat Anti Mouse IgG IgA IgM (STAR87...) HRP

Recommended Negative Controls

MOUSE IgM NEGATIVE CONTROL (MCA692)

North & South Tel: +1 800 265 7376 America Fax: +1 919 878 3751 Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

Printed on 28 Jan 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint