

Datasheet: MCA4610T

BATCH NUMBER 156536

Description:	MOUSE ANTI HUMAN CD15
Specificity:	CD15
Other names:	LEWIS X
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	HI98
Isotype:	IgM
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/5 - 1/10
Immunohistology - Frozen	▪			
Immunohistology - Paraffin	▪			
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species

Human

Species Cross Reactivity

Reacts with: Chimpanzee

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgM - liquid

Preparation	Ig fraction prepared by affinity chromatography
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	Ig concentration 0.5mg/ml
RRID	AB_2247102
Specificity	Mouse anti Human CD15 antibody, clone HI98 recognises human CD15, otherwise known as Lewis X and stage-specific embryonic antigen-1 (SSEA-1). The CD15 antigen is itself a terminal trisaccharide, Gal-beta1-4 (Fuc-alpha1-3) GlcNAc (3-FAL), which is found on a variety of glycoproteins and glycolipids. CD15 is predominately expressed by peripheral blood granulocytes and various epithelial tissues. It plays an important role in cell adhesion and has been used as a marker for diagnosis of Hodgkin's lymphoma (Zukerberg <i>et al.</i> 1991).
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.
Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	Guaranteed until date of expiry. Please see product label.
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA4610T 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgM (STAR138...) [Alk. Phos.](#)

Goat Anti Mouse IgG IgA IgM (STAR87...) [HRP](#)

Recommended Negative Controls

[MOUSE IgM NEGATIVE CONTROL \(MCA692\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)