Datasheet: MCA461F BATCH NUMBER 1806

Description:	MOUSE ANTI HUMAN CD45RO:FITC
Specificity:	CD45RO
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	UCHL1
lsotype:	lgG2a
Quantity:	0.1 mg

Product Details

Applications								
derived from testing within our laboratories, peer-reviewed publications or pe communications from the originators. Please refer to references indicated for								
	information. For general protocol recommendations, please visit <u>www.bio-</u>							
	-	rad-antibodies.com/protocols.						
		Yes	No	Not Determined	Suggested Dilution			
	Flow Cytometry	-			Neat			
Where this antibody has not been tested for use in a particular technique this does necessarily exclude its use in such procedures. Suggested working dilutions are gi a guide only. It is recommended that the user titrates the antibody for use in their of system using appropriate negative/positive controls.								
Target Species	Human							
Species Cross Reactivity	Reacts with: Chimpanzee, Marmoset, Pig N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.							
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid							
Max Ex/Em	Fluorophore FITC	Excitation Ma 490	x (nm)	Emission Max (nm) 525				
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant							
Buffer Solution	Phosphate buffered sa	line						

Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
Immunogen	Human IL-2 dependent T cells
External Database Links	UniProt: <u>P08575</u> <u>Related reagents</u> Entrez Gene: <u>5788</u> PTPRC <u>Related reagents</u>
Synonyms	CD45
RRID	AB_321433
Fusion Partners	Spleen cells from immunised mice were fused with cells of the mouse P3/NS1/1-Ag4-1 myeloma cell line
Specificity	 Mouse anti Human CD45RO monoclonal antibody, clone UCHL1 recognizes the low molecular weight isoform (180 kDa) of the leucocyte common antigen (LCA). The antigen is expressed by a functional subset of T cells with memory phenotype. In peripheral blood Mouse anti Human CD45RO, clone UCHL1 stains 40-80% of lymphocytes and all monocytes and granulocytes.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells or 100ul whole blood
References	 Smith, S.H. <i>et al.</i> (1986) Functional subsets of human helper-inducer cells defined by a new monoclonal antibody, UCHL1. <u>Immunology 58: 63-70.</u> Norton, A.J. <i>et al.</i> (1986) Monoclonal antibody (UCHL1) that recognises normal and neoplastic T cells in routinely fixed tissues. <u>J Clin Pathol. 39 (4): 399-405.</u> Beverley, P.C.L. <i>et al.</i> (1986) T-cell subsets and function. Progress in Immunology VI. Cinader, B., Miller, G.G., eds. Academic Press Orlando pp 941-948. Beverley, P.C. (1987) Human T cell subsets. <u>Immunol Lett. 14 (4): 263-7.</u> Terry. L.A. <i>et al.</i> (1987) Phenotypic heterogeneity of the CD4+ and CD8+ subsets. Leucocyte Typing III. McMichael, A.J., Beverley, P.C.L. <i>et al.</i> eds. University Press. pp 225-7. Akbar, A.N. <i>et al.</i> (1988) Loss of CD45R and gain of UCHL1 reactivity is a feature of primed T cells. <u>J Immunol. 140 (7): 2171-8.</u> Terry, L.A. <i>et al.</i> (1988) The monoclonal antibody, UCHL1, recognizes a 180,000 MW component of the human leucocyte-common antigen, CD45. <u>Immunology. 64 (2): 331-6.</u> Beverley, P.C.L. <i>et al.</i> (1988) Phenotypic diversity of the CD45 antigen and its relation to function. <u>Immunology, Suppl. 1: 3-5.</u> Merkenschlager, M. <i>et al.</i> (1988) Limiting dilution analysis of proliferative responses in

	 human lymphocyte populations defined by the monoclonal antibody UCHL1: implications for differential CD45 expression in T cell memory formation. <u>Eur J Immunol. 18 (11):</u> 1653-61. 10. Cavers, M. <i>et al.</i> (2002) Differential expression of beta1 and beta2 integrins and L-selectin on CD4+ and CD8+ T lymphocytes in human blood: comparative analysis between isolated cells, whole blood samples and cryopreserved preparations. <u>Clin Exp Immunol. 127: 60-5.</u> 11. Hutnick, N.A. <i>et al.</i> (2010) Vaccination with Ad5 vectors expands Ad5-specific CD8 T cells without altering memory phenotype or functionality. <u>PLoS One. 5: e14385.</u> 12. Leigh, J.E. <i>et al.</i> (2006) Characterization of the immune status of CD8+ T cells in oral lesions of human immunodeficiency virus-infected persons with oropharyngeal Candidiasis. <u>Clin Vaccine Immunol. 13: 678-83.</u> 13. Nistala, K. <i>et al.</i> (2008) Interleukin-17-producing T cells are enriched in the joints of children with arthritis, but have a reciprocal relationship to regulatory T cell numbers. <u>Arthritis Rheum. 58: 875-87.</u> 14. Liu Y <i>et al.</i> (2015) Fractionation of human spermatogenic cells using STA-PUT gravity sedimentation and their miRNA profiling. <u>Sci Rep. 5: 8084.</u> 15. Wilson, C.L. <i>et al.</i> (2015) Ubiquitin C-terminal hydrolase 1: A novel functional marker for liver myofibroblasts and a therapeutic target in chronic liver disease. <u>J Hepatol. 63 (6):</u> 1421-8. 16. Zhao, H. <i>et al.</i> (2018) <i>In vitro</i> differentiation of spermatogonial stem cells using testicular cells from Guangxi Bama mini-pig. <u>J Vet Sci. 19 (5): 592-9.</u>
Storage	Store at +4°C or at -20°C if preferred.
	This product should be stored undiluted.
	Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.
	Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA461F 10041
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG2a NEGATIVE CONTROL:FITC (MCA929F)

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio	-rad.com	Email: antibody_sales_uk@bio	o-rad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M367720:200529'

Printed on 08 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint