

## Datasheet: MCA4633GA

<b>Description:</b>	MOUSE ANTI RAT CD106
<b>Specificity:</b>	CD106
<b>Other names:</b>	VCAM-1
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	MR106
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/25
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	
Immunofluorescence	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Rat
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> )

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
Immunogen	Rat CD106-transfected L5178Y cells.
External Database Links	<p><b>UniProt:</b>  <a href="#">P29534</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">25361</a>    Vcam1    <a href="#">Related reagents</a></p>
Synonyms	Vcam-1
RRID	AB_1604705
Specificity	<p><b>Mouse anti Rat CD106 antibody, clone MR106</b> recognizes rat CD106, otherwise known as VCAM-1 (vascular adhesion molecule 1), a 110kDa inducible type I transmembrane glycoprotein and member of the immunoglobulin supergene family, which is predominantly expressed on vascular endothelium, and also on bone marrow stromal cells, follicular dendritic cells and some macrophages.</p> <p>CD106 interacts with the lymphocyte homing receptor VLA-4 (alpha4 beta1 integrin) and LPAM-1 (alpha4 beta7 integrin) and mediates leucocyte-endothelial cell adhesion and signal transduction. Expression of CD106 is upregulated following cytokine activation, and endothelial CD106 plays a role in the extravasation of leucocytes from blood vessels during inflammation.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label $1 \times 10^6$ cells in 100ul.
References	<p>1. Miyao, N. <i>et al.</i> (2006) Various adhesion molecules impair microvascular leukocyte kinetics in ventilator-induced lung injury. <a href="#">Am J Physiol Lung Cell Mol Physiol. 290 (6): L1059-68.</a></p> <p>2. Kubota, H. <i>et al.</i> (2007) Identification and characterization of vitamin A-storing cells in fetal liver: implications for functional importance of hepatic stellate cells in liver development and hematopoiesis. <a href="#">Stem Cells. 25 (9): 2339-49.</a></p>
Further Reading	1. Bevilacqua, M.P. (1993) Endothelial-leukocyte adhesion molecules. <a href="#">Annu Rev Immunol. 11: 767-804.</a>
Storage	<p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.</p>

<b>Guarantee</b>	12 months from date of despatch
<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA4633GA">https://www.bio-rad-antibodies.com/SDS/MCA4633GA</a> 10040
<b>Regulatory</b>	For research purposes only

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	<a href="#">RPE</a>
Goat Anti Mouse IgG IgA IgM (STAR87...)	<a href="#">Alk. Phos.</a> , <a href="#">HRP</a>
Goat Anti Mouse IgG (STAR76...)	<a href="#">RPE</a>
Rabbit Anti Mouse IgG (STAR13...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (STAR70...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (H/L) (STAR117...)	<a href="#">Alk. Phos.</a> , <a href="#">DyLight®488</a> , <a href="#">DyLight®550</a> , <a href="#">DyLight®650</a> , <a href="#">DyLight®680</a> , <a href="#">DyLight®800</a> , <a href="#">FITC</a> , <a href="#">HRP</a>
Goat Anti Mouse IgG (Fc) (STAR120...)	<a href="#">FITC</a> , <a href="#">HRP</a>
Goat Anti Mouse IgG (STAR77...)	<a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR9...)	<a href="#">FITC</a>

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA1209\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)  
'M384161:210513'

Printed on 18 Jan 2024