

### Datasheet: MCA4633PE

Description:	MOUSE ANTI RAT CD106:RPE
Specificity:	CD106
Other names:	VCAM-1
Format:	RPE
Product Type:	Monoclonal Antibody
Product Type: Clone:	Monoclonal Antibody MR106
	•

## **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	-			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Rat					
Product Form	Purified IgG conjuga	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized				
Reconstitution	Reconstitute with 1r	Reconstitute with 1ml distilled water				
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)			
	RPE 488nm laser	496	578			
Preparation	Purified IgG prepare supernatant	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant				
Buffer Solution	Phosphate buffered	saline				
Preservative Stabilisers	0.09% Sodium Azid 1% Bovine Serum A 5% Sucrose	, ,,				

Immunogen	Rat CD106-transfected L5178Y cells.
External Database Links	UniProt: P29534 Related reagents  Entrez Gene: 25361 Vcam1 Related reagents
Synonyms	Vcam-1
RRID	AB_2214213
Specificity	Mouse anti Rat CD106 antibody, clone MR106 recognizes rat CD106, otherwise known as VCAM-1 (vascular adhesion molecule 1), a 110kDa inducible type I transmembrane glycoprotein and member of the immunoglobulin supergene family, which is predominantly expressed on vascular endothelium, and also on bone marrow stromal cells, follicular dendritic cells and some macrophages.
	CD106 interacts with the lymphocyte homing receptor VLA-4 (alpha4 beta1 integrin) and LPAM-1 (alpha4 beta7 integrin) and mediates leucocyte-endothelial cell adhesion and signal transduction. Expression of CD106 is upregulated following cytokine activation, and endothelial CD106 plays a role in the extravasation of leucocytes from blood vessels during inflammation.
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul.
References	<ol> <li>Miyao, N. <i>et al.</i> (2006) Various adhesion molecules impair microvascular leukocyte kinetics in ventilator-induced lung injury. <u>Am J Physiol Lung Cell Mol Physiol. 290 (6): L1059-68.</u></li> <li>Kubota, H. <i>et al.</i> (2007) Identification and characterization of vitamin A-storing cells in fetal liver: implications for functional importance of hepatic stellate cells in liver development and hematopoiesis. <u>Stem Cells. 25 (9): 2339-49.</u></li> </ol>
Further Reading	1. Bevilacqua, M.P. (1993) Endothelial-leukocyte adhesion molecules. <u>Annu Rev Immunol.</u> 11: 767-804.
Storage	Prior to reconstitution store at +4°C.  After reconstitution store at +4°C.  DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #20487 available at: 20487: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/20487.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/20487.pdf</a>
Regulatory	For research purposes only

# **Related Products**

## **Recommended Negative Controls**

### MOUSE IgG1 NEGATIVE CONTROL:RPE (MCA1209PE)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 **Europe** Fax: +44 (0)1865 852 739

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

 Email: antibody\_sales\_de@bio-rad.com

'M375546:210104'

### Printed on 04 Jan 2021

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint