

Datasheet: MCA4635T

BATCH NUMBER 1612

Description:	RAT ANTI MOUSE CD4
Specificity:	CD4
Other names:	L3T4 ANTIGEN, LY-4
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	GK1.5
Isotype:	IgG2b
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/200
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA	▪			
Immunoprecipitation	▪			
Western Blotting	▪			
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative	0.09% Sodium Azide

Stabilisers

Carrier Free Yes

Approx. Protein Concentrations IgG concentration 1.0 mg/ml

Immunogen Murine CD4.

External Database Links

UniProt:
[P06332](#) [Related reagents](#)

Entrez Gene:
[12504](#) Cd4 [Related reagents](#)

RRID AB_1898234

Fusion Partners Spleen cells from immunised Lewis rats were fused with cells of the SP2/0 myeloma cell line.

Specificity

Rat anti Mouse CD4 antibody, clone GK1.5 recognizes mouse CD4, a ~55 kDa protein also known as Ly-4 and L3T4. CD4 is a single chain transmembraneous glycoprotein which belongs to the immunoglobulin superfamily, and is primarily expressed on T helper cells, peripheral blood monocytes and tissue macrophages. CD4 is also expressed on a subpopulation of regulatory T cells (CD4⁺ CD25⁺), which play a key role in the maintenance of self tolerance.

Rat anti Mouse CD4 antibody, clone GK1.5 has been reported to block CD4⁺ T-cell activation. It blocks class II MHC antigen-specific binding, thereby inhibiting functions such as class II MHC antigen-specific proliferation and the release of lymphokines. It may also be used for *in vivo* and *in vitro* cell depletion of CD4⁺ T-cells.

Flow Cytometry Use 10ul of the suggested working dilution to label 1x10⁶ cells in 100ul.

References

1. Dialynas, D.P. *et al.* (1983) Characterization of the murine T cell surface molecule, designated L3T4, identified by monoclonal antibody GK1.5: similarity of L3T4 to the human Leu-3/T4 molecule. [J Immunol. 131 \(5\): 2445-51.](#)
2. Wilde, D.B. *et al.* (1983) Evidence implicating L3T4 in class II MHC antigen reactivity; monoclonal antibody GK1.5 (anti-L3T4a) blocks class II MHC antigen-specific proliferation, release of lymphokines, and binding by cloned murine helper T lymphocyte lines. [J Immunol. 131 \(5\): 2178-83.](#)
3. Dialynas, D.P. *et al.* (1983) Characterization of the murine antigenic determinant, designated L3T4a, recognized by monoclonal antibody GK1.5: expression of L3T4a by functional T cell clones appears to correlate primarily with class II MHC antigen-reactivity. [Immunol Rev. 74: 29-56.](#)
4. Zhou, Z. *et al.* (2011) Autoreactive marginal zone B cells enter the follicles and interact with CD4⁺ T cells in lupus-prone mice. [BMC Immunol. 12: 7.](#)
5. Näher, H. *et al.* (1985) Dynamics of T cells of L3T4 and Ly 2 phenotype within

- granulomas in murine listeriosis. [Clin Exp Immunol. 60 \(3\): 559-64.](#)
6. Ye, X. *et al.* (2000) Transient depletion of CD4 lymphocyte improves efficacy of repeated administration of recombinant adenovirus in the ornithine transcarbamylase deficient sparse fur mouse. [Gene Ther. 7 \(20\): 1761-7.](#)
7. Chu, N.R. *et al.* (2000) Immunotherapy of a human papillomavirus (HPV) type 16 E7-expressing tumour by administration of fusion protein comprising *Mycobacterium bovis* bacille Calmette-Guérin (BCG) hsp65 and HPV16 E7. [Clin Exp Immunol. 121:216-25](#)
8. Pletinckx, K. *et al.* (2015) Immature dendritic cells convert anergic nonregulatory T cells into Foxp3- IL-10+ regulatory T cells by engaging CD28 and CTLA-4. [Eur J Immunol. 45 \(2\): 480-91.](#)
9. Foy, S.P. *et al.* (2016) Poxvirus-Based Active Immunotherapy with PD-1 and LAG-3 Dual Immune Checkpoint Inhibition Overcomes Compensatory Immune Regulation, Yielding Complete Tumor Regression in Mice. [PLoS One. 11 \(2\): e0150084.](#)
10. Steinl, D.C. *et al.* (2016) Noninvasive Contrast-Enhanced Ultrasound Molecular Imaging Detects Myocardial Inflammatory Response in Autoimmune Myocarditis. [Circ Cardiovasc Imaging. 9 \(8\): .](#)
11. Olesen, M. N. *et al.* (2018) CD4 T cells react to local increase of α -synuclein in a pathology-associated variant-dependent manner and modify brain microglia in absence of brain pathology [Heliyon. 4 \(1\): e00513.](#)

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA4635T>
10040

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR73...)	RPE
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight@550 , DyLight@650 , DyLight@800
Rabbit Anti Rat IgG (STAR21...)	HRP
Rabbit Anti Rat IgG (STAR16...)	DyLight@800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (STAR72...)	HRP

North & South Tel: +1 800 265 7376

Worldwide Tel: +44 (0)1865 852 700

Europe Tel: +49 (0) 89 8090 95 21

America Fax: +1 919 878 3751

Fax: +44 (0)1865 852 739

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To
find a

batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M367754:200529'

Printed on 12 Aug 2023

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)