

Datasheet: MCA4635T

Description:	RAT ANTI MOUSE CD4
Specificity:	CD4
Other names:	L3T4 ANTIGEN, LY-4
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	GK1.5
Isotype:	IgG2b
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/200
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA	▪			
Immunoprecipitation	▪			
Western Blotting	▪			
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Murine CD4.
External Database Links	<p>UniProt: P06332 Related reagents</p> <p>Entrez Gene: 12504 Cd4 Related reagents</p>
RRID	AB_1898234
Fusion Partners	Spleen cells from immunised Lewis rats were fused with cells of the SP2/0 myeloma cell line.
Specificity	<p>Rat anti Mouse CD4 antibody, clone GK1.5 recognizes mouse CD4, a ~55 kDa protein also known as Ly-4 and L3T4. CD4 is a single chain transmembraneous glycoprotein which belongs to the immunoglobulin superfamily, and is primarily expressed on T helper cells, peripheral blood monocytes and tissue macrophages. CD4 is also expressed on a subpopulation of regulatory T cells (CD4⁺ CD25⁺), which play a key role in the maintenance of self tolerance.</p> <p>Rat anti Mouse CD4 antibody, clone GK1.5 has been reported to block CD4⁺ T-cell activation. It blocks class II MHC antigen-specific binding, thereby inhibiting functions such as class II MHC antigen-specific proliferation and the release of lymphokines. It may also be used for <i>in vivo</i> and <i>in vitro</i> cell depletion of CD4⁺ T-cells.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> Dialynas, D.P. <i>et al.</i> (1983) Characterization of the murine T cell surface molecule, designated L3T4, identified by monoclonal antibody GK1.5: similarity of L3T4 to the human Leu-3/T4 molecule. J Immunol. 131 (5): 2445-51. Wilde, D.B. <i>et al.</i> (1983) Evidence implicating L3T4 in class II MHC antigen reactivity; monoclonal antibody GK1.5 (anti-L3T4a) blocks class II MHC antigen-specific proliferation, release of lymphokines, and binding by cloned murine helper T lymphocyte lines. J Immunol. 131 (5): 2178-83. Näher, H. <i>et al.</i> (1985) Dynamics of T cells of L3T4 and Ly 2 phenotype within granulomas in murine listeriosis. Clin Exp Immunol. 60 (3): 559-64. Ye, X. <i>et al.</i> (2000) Transient depletion of CD4 lymphocyte improves efficacy of repeated administration of recombinant adenovirus in the ornithine transcarbamylase deficient sparse fur mouse. Gene Ther. 7 (20): 1761-7. Chu, N.R. <i>et al.</i> (2000) Immunotherapy of a human papillomavirus (HPV) type 16 E7-expressing tumour by administration of fusion protein comprising <i>Mycobacterium bovis</i> bacille Calmette-Guérin (BCG) hsp65 and HPV16 E7. Clin Exp Immunol. 121:216-25

6. Zhou, Z. *et al.* (2011) Autoreactive marginal zone B cells enter the follicles and interact with CD4+ T cells in lupus-prone mice. [BMC Immunol. 12: 7.](#)
7. Pletinckx, K. *et al.* (2015) Immature dendritic cells convert anergic nonregulatory T cells into Foxp3- IL-10+ regulatory T cells by engaging CD28 and CTLA-4. [Eur J Immunol. 45 \(2\): 480-91.](#)
8. Foy, S.P. *et al.* (2016) Poxvirus-Based Active Immunotherapy with PD-1 and LAG-3 Dual Immune Checkpoint Inhibition Overcomes Compensatory Immune Regulation, Yielding Complete Tumor Regression in Mice. [PLoS One. 11 \(2\): e0150084.](#)
9. Steinl, D.C. *et al.* (2016) Noninvasive Contrast-Enhanced Ultrasound Molecular Imaging Detects Myocardial Inflammatory Response in Autoimmune Myocarditis. [Circ Cardiovasc Imaging. 9 \(8\): .](#)
10. Olesen, M. N. *et al.* (2018) CD4 T cells react to local increase of α -synuclein in a pathology-associated variant-dependent manner and modify brain microglia in absence of brain pathology [Heliyon. 4 \(1\): e00513.](#)
11. Jalili, R.B. *et al.* (2018) Fibroblast cell-based therapy prevents induction of alopecia areata in an experimental model. [Cell Transplant. 27 \(6\): 994-1004.](#)
12. Zheng, Z. *et al.* (2022) *In Vivo*. Inhibition of TRPC6 by SH045 Attenuates Renal Fibrosis in a New Zealand Obese (NZO) Mouse Model of Metabolic Syndrome. [Int J Mol Sci. 23\(12\):6870.](#)

Further Reading 1. Dialynas, D.P. *et al.* (1983) Characterization of the murine antigenic determinant, designated L3T4a, recognized by monoclonal antibody GK1.5: expression of L3T4a by functional T cell clones appears to correlate primarily with class II MHC antigen-reactivity. [Immunol Rev. 74: 29-56.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA4635T>
10040

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Rat IgG (STAR69...) [FITC](#)
 Goat Anti Rat IgG (STAR73...) [RPE](#)
 Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...) [DyLight@550](#), [DyLight@650](#), [DyLight@800](#)
 Rabbit Anti Rat IgG (STAR21...) [HRP](#)

Rabbit Anti Rat IgG (STAR16...)

[DyLight®800](#)

Goat Anti Rat IgG (STAR131...)

[Alk. Phos., Biotin](#)

Rabbit Anti Rat IgG (STAR17...)

[FITC](#)

Goat Anti Rat IgG (STAR72...)

[HRP](#)

North & South Tel: +1 800 265 7376

Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21

America Fax: +1 919 878 3751

Fax: +44 (0)1865 852 739

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M384162:210513'

Printed on 12 Aug 2023

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)