

## Datasheet: MCA4635T

<b>Description:</b>	RAT ANTI MOUSE CD4
<b>Specificity:</b>	CD4
<b>Other names:</b>	L3T4 ANTIGEN, LY-4
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	GK1.5
<b>Isotype:</b>	IgG2b
<b>Quantity:</b>	25 µg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/200
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA	▪			
Immunoprecipitation	▪			
Western Blotting	▪			
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Mouse
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide
<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Murine CD4.

**External Database  
Links**

**UniProt:**

[P06332](#) [Related reagents](#)

**Entrez Gene:**

[12504](#) Cd4 [Related reagents](#)

---

**Fusion Partners**

Spleen cells from immunised Lewis rats were fused with cells of the SP2/0 myeloma cell line.

---

**Specificity**

**Rat anti Mouse CD4 antibody, clone GK1.5** recognizes mouse CD4, a ~55 kDa protein also known as Ly-4 and L3T4. CD4 is a single chain transmembrane glycoprotein which belongs to the immunoglobulin superfamily, and is primarily expressed on T helper cells, peripheral blood monocytes and tissue macrophages. CD4 is also expressed on a subpopulation of regulatory T cells (CD4<sup>+</sup> CD25<sup>+</sup>), which play a key role in the maintenance of self tolerance.

Rat anti Mouse CD4 antibody, clone GK1.5 has been reported to block CD4<sup>+</sup> T-cell activation. It blocks class II MHC antigen-specific binding, thereby inhibiting functions such as class II MHC antigen-specific proliferation and the release of lymphokines. It may also be used for *in vivo* and *in vitro* cell depletion of CD4<sup>+</sup> T-cells.

---

**Flow Cytometry**

Use 10ul of the suggested working dilution to label 1x10<sup>6</sup> cells in 100ul.

---

**References**

1. Dialynas, D.P. *et al.* (1983) Characterization of the murine T cell surface molecule, designated L3T4, identified by monoclonal antibody GK1.5: similarity of L3T4 to the human Leu-3/T4 molecule. [J Immunol. 131 \(5\): 2445-51.](#)
2. Wilde, D.B. *et al.* (1983) Evidence implicating L3T4 in class II MHC antigen reactivity; monoclonal antibody GK1.5 (anti-L3T4a) blocks class II MHC antigen-specific proliferation, release of lymphokines, and binding by cloned murine helper T lymphocyte lines. [J Immunol. 131 \(5\): 2178-83.](#)
3. Dialynas, D.P. *et al.* (1983) Characterization of the murine antigenic determinant, designated L3T4a, recognized by monoclonal antibody GK1.5: expression of L3T4a by functional T cell clones appears to correlate primarily with class II MHC antigen-reactivity. [Immunol Rev. 74: 29-56.](#)
4. Zhou, Z. *et al.* (2011) Autoreactive marginal zone B cells enter the follicles and interact with CD4<sup>+</sup> T cells in lupus-prone mice. [BMC Immunol. 12: 7.](#)
5. Näher, H. *et al.* (1985) Dynamics of T cells of L3T4 and Ly 2 phenotype within granulomas in murine listeriosis. [Clin Exp Immunol. 60 \(3\): 559-64.](#)
6. Ye, X. *et al.* (2000) Transient depletion of CD4 lymphocyte improves efficacy of repeated administration of recombinant adenovirus in the ornithine transcarbamylase deficient sparse fur mouse. [Gene Ther. 7 \(20\): 1761-7.](#)
7. Chu, N.R. *et al.* (2000) Immunotherapy of a human papillomavirus (HPV) type 16 E7-expressing tumour by administration of fusion protein comprising *Mycobacterium bovis* bacille Calmette-Guérin (BCG) hsp65 and HPV16 E7. [Clin Exp Immunol. 121:216-25](#)
8. Pletinckx, K. *et al.* (2015) Immature dendritic cells convert anergic nonregulatory T cells into Foxp3- IL-10<sup>+</sup> regulatory T cells by engaging CD28 and CTLA-4. [Eur J Immunol. 45 \(2\): 480-91.](#)
9. Foy, S.P. *et al.* (2016) Poxvirus-Based Active Immunotherapy with PD-1 and LAG-3 Dual Immune Checkpoint Inhibition Overcomes Compensatory Immune Regulation, Yielding Complete Tumor Regression in Mice. [PLoS One. 11 \(2\): e0150084.](#)
10. Steinl, D.C. *et al.* (2016) Noninvasive Contrast-Enhanced Ultrasound Molecular Imaging Detects Myocardial Inflammatory Response in Autoimmune Myocarditis. [Circ Cardiovasc Imaging. 9 \(8\): .](#)
11. Olesen, M. N. *et al.* (2018) CD4 T cells react to local increase of α-synuclein in a pathology-associated variant-dependent manner and modify brain microglia in absence of brain pathology [Heliyon. 4 \(1\): e00513.](#)

<b>Storage</b>	Store at +4°C or at -20°C if preferred.  This product should be stored undiluted.  Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
<b>Shelf Life</b>	18 months from date of despatch.
<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10040 available at: 10040: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf</a>
<b>Regulatory</b>	For research purposes only

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	<a href="#">DyLight®800</a>
Rabbit Anti Rat IgG (STAR17...)	<a href="#">FITC</a>
Goat Anti Rat IgG (STAR69...)	<a href="#">FITC</a>
Goat Anti Rat IgG (STAR72...)	<a href="#">HRP</a>
Goat Anti Rat IgG (STAR131...)	<a href="#">Alk. Phos.</a> , <a href="#">Biotin</a>
Goat Anti Rat IgG (STAR73...)	<a href="#">RPE</a>
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	<a href="#">DyLight®649</a> , <a href="#">DyLight®800</a>
Rabbit Anti Rat IgG (STAR21...)	<a href="#">HRP</a>

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

'M319155:180719'

Printed on 18 Sep 2018