

## Datasheet: MCA4636

<b>Description:</b>	HAMSTER ANTI MOUSE CD152
<b>Specificity:</b>	CD152
<b>Other names:</b>	CTLA-4
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	1B8
<b>Isotype:</b>	IgG
<b>Quantity:</b>	0.25 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/100
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			
Immunoprecipitation	▪			
Western Blotting	▪			
Immunofluorescence	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Mouse
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> )

<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0mg/ml
<b>Immunogen</b>	Extracellular portion of mouse CD152 fused to a mouse IgG2a.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P09793</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">12477</a>    Ctla4    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	Cd152
<b>RRID</b>	AB_10547302
<b>Fusion Partners</b>	Lymph cells from immunised Armenian hamsters were fused with cells of the murine SP2/0 myeloma cell line.
<b>Specificity</b>	<p><b>Hamster anti Mouse CD152 antibody, clone 1B8</b> recognizes mouse CD152, commonly known as CTLA-4 (cytotoxic T-lymphocyte protein 4), a structural homolog for CD28. CD152 binds the same ligands as CD28, namely CD80 (B7.1) and CD86 (B7.2), but with much higher affinity. However, while binding of CD28 enhances T cell function, binding of CD152 inhibits it. CD152 is not detected on naïve T-cells and is induced upon T-cell activation.</p> <p>In mice, lack of CD152 leads to massive lymphoproliferative disorder leading to tissue destruction and autoimmunity, suggesting a critical role for CD152 in limiting T-cell responses. CD152 has been shown to reduce IL-2 production after ligation and can inhibit T-cell expansion. This regulatory role and the high binding affinity of CD152 make it a potential therapy for autoimmune diseases such as rheumatoid arthritis.</p> <p>Hamster anti Mouse CD152 antibody, clone 1B8 may be used to enhance proliferation and pro-inflammatory cytokines of lymph node cells. It may also be used to enhance tumour immunity.</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label $1 \times 10^6$ cells in 100ul.
<b>References</b>	<ol style="list-style-type: none"> <li>Walunas, T. L. <i>et al.</i> (1994) CTLA-4 can function as a negative regulator of T cell activation. <a href="#">Immunity 1: 405-413.</a></li> <li>Verhagen, J. <i>et al.</i> (2013) CTLA-4 controls the thymic development of both conventional and regulatory T cells through modulation of the TCR repertoire. <a href="#">Proc Natl Acad Sci U S A. 110: E221-30.</a></li> </ol>
<b>Further Reading</b>	1. Karandikar, N.J. <i>et al.</i> (1996) CTLA-4: a negative regulator of autoimmune disease. <a href="#">J Exp Med. 184 (2): 783-8.</a>

**Storage** This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

---

**Guarantee** 12 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA4636>  
10040

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Goat Anti Hamster IgG (STAR104...) [DyLight@800](#), [FITC](#)

Goat Anti Hamster IgG (STAR79...) [Biotin](#), [FITC](#), [HRP](#)

### Recommended Negative Controls

[HAMSTER \(ARMENIAN\) IgG NEGATIVE CONTROL \(MCA2356\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M384163:210513'

**Printed on 12 Aug 2023**

---

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)