

Datasheet: MCA4636GA

Description:	HAMSTER ANTI MOUSE CD152
Specificity:	CD152
Other names:	CTLA-4
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	1B8
Isotype:	IgG
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			1/10 - 1/100
Immunohistology - Frozen				
Immunohistology - Paraffin			•	
ELISA	•			
Immunoprecipitation	•			
Western Blotting	•			
Immunofluorescence	•			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse	
Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared by affinity chromatography on Protein G supernatant	From tissue culture
Buffer Solution	Phosphate buffered saline	
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)	

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
Immunogen	Extracellular portion of mouse CD152 fused to a mouse IgG2a.
External Database Links	UniProt: P09793 Related reagents
	Entrez Gene: 12477 Ctla4 Related reagents
Synonyms	Cd152
RRID	AB_2087650
Fusion Partners	Lymph cells from immunized Armenian hamsters were fused with cells of the murine SP2/0 myeloma cell line.
Specificity	Hamster anti Mouse CD152 antibody, clone 1B8 recognizes mouse CD152, also known as CTLA-4 (cytotoxic T-lymphocyte protein 4), a structural homolog for CD28. CD152 binds the same ligands as CD28, namely CD80 (B7.1) and CD86 (B7.2), but with much higher affinity. However, while binding of CD28 enhances T cell function, binding of CD152 inhibits it. CD152 is not detected on naïve T-cells and is induced upon T-cell activation. In mice, lack of CD152 leads to massive lymphoproliferative disorder leading to tissue
	destruction and autoimmunity, suggesting a critical role for CD152 in limiting T-cell responses. CD152 has been shown to reduce IL-2 production after ligation and can inhibit T-cell expansion. This regulatory role and the high binding affinity of CD152 make it a potential therapy for autoimmune diseases such as rheumatoid arthritis.
	Hamster anti Mouse CD152 antibody, clone 1B8 may be used to enhance proliferation and pro-inflammatory cytokines of lymph node cells. It may also be used to enhance tumor immunity.
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.
References	 Walunas, T. L. <i>et al.</i> (1994) CTLA-4 can function as a negative regulator of T cell activation. <u>Immunity 1: 405-413.</u> Verhagen, J. <i>et al.</i> (2013) CTLA-4 controls the thymic development of both conventional and regulatory T cells through modulation of the TCR repertoire. <u>Proc Natl Acad Sci U S A. 110 (3): E221-30.</u>
Further Reading	1. Karandikar, N.J. <i>et al.</i> (1996) CTLA-4: a negative regulator of autoimmune disease. <u>J</u> Exp Med. 184 (2): 783-8.
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at

-20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA4636GA 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Hamster IgG (STAR104...) <u>DyLight®550</u>, <u>DyLight®650</u>, <u>DyLight®800</u>,

FITC

Goat Anti Hamster IgG (STAR79...) Biotin, FITC, HRP

Recommended Negative Controls

HAMSTER (ARMENIAN) IgG NEGATIVE CONTROL (MCA2356)

North & South Tel: +1 800 265 7376 Worldwide Tel: +44 (0)1865 852 700

America Fax: +1 919 878 3751 Fax: +44 (0)1865 852 739

 $\textbf{Email: antibody_sales_us@bio-rad.com} \\ \textbf{Email: antibody_sales_uk@bio-rad.com} \\ \textbf{Email: antibody_sales_de@bio-rad.com} \\ \textbf{Email: antibody_sales_de@b$

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

2 2 3

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M384509:210513'

Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint