

Datasheet: MCA4643FT

| Description:         | MOUSE ANTI HUMAN CD96:FITC |
|----------------------|----------------------------|
| Specificity:         | CD96                       |
| Other names:         | TACTILE                    |
| Format:              | FITC                       |
| <b>Product Type:</b> | Monoclonal Antibody        |
| Clone:               | NK92.39                    |
| Isotype:             | IgG1                       |
| Quantity:            | 25 μg                      |
|                      |                            |

#### **Product Details**

### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

|                | Yes | No | Not Determined | Suggested Dilution |
|----------------|-----|----|----------------|--------------------|
| Flow Cytometry | •   |    |                | Neat - 1/10        |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

| Target Species                    | Human                             |                         |                     |
|-----------------------------------|-----------------------------------|-------------------------|---------------------|
| Product Form                      | Purified IgG conjugate            | ed to Fluorescein Isoth | niocyanate Isomer 1 |
| Max Ex/Em                         | Fluorophore                       | Excitation Max (nm)     | Emission Max (nm    |
|                                   | FITC                              | 490                     | 525                 |
| Preparation                       | Purified IgG prepared supernatant |                         | raphy on Protein G  |
| Buffer Solution                   | Phosphate buffered s              | aline                   |                     |
| Preservative                      | 0.09% Sodium Azide                | (NaN <sub>3</sub> )     |                     |
| Stabilisers                       | 1% Bovine Serum Alb               | oumin                   |                     |
| Approx. Protein<br>Concentrations | IgG concentration 0.1             | mg/ml                   |                     |

| Immunogen                        | Natural killer 92 (NK92) cell line.   |
|----------------------------------|---|
| External Database<br>Links       | UniProt:  P40200 Related reagents  Entrez Gene:  10225 CD96 Related reagents  |
| RRID                             | AB_1628793  |
| Specificity                      | Mouse anti Human CD96 antibody, clone NK92.39 recognizes human CD96, otherwise known as Tactile (T cell-activated increased late expression), a type I transmembrane glycoprotein expressed at low levels on peripheral resting T and NK cells, which increases following activation.   |
|                                  | CD96 stimulates the cytotoxicity of activated NK cells, and promotes the adhesion of NK cells to target cells expressing CD155/poliovirus receptor ( <u>PVR</u> ) and the transfer of target cell PVR to the NK cells. CD96 can be used in the immunophenotyping of T-cell acute lymphoblastic leukaemias ( <u>ALL</u> ) and acute myeloid leukaemias ( <u>AML</u> ), and may play a role in the recognition of CD155-expressing tumours by NK cells. |
|                                  | Mouse anti Human CD96 antibody, clone NK92.39 blocks the binding of soluble CD155 to NK92 cells ( <u>Fuchs <i>et al.</i> 2004</u> ).  |
| Flow Cytometry                   | Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul.   |
| References                       | 1. Fuchs, A. <i>et al.</i> (2004) Cutting edge: CD96 (tactile) promotes NK cell-target cell adhesion by interacting with the poliovirus receptor (CD155). <u>J Immunol. 172 (7): 3994-8.</u>  |
| Storage                          | Store at +4°C or at -20°C if preferred.  Storage in frost-free freezers is not recommended.  This product should be stored undiluted. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.  |
| Guarantee                        | 12 months from date of despatch   |
| Health And Safety<br>Information | Material Safety Datasheet documentation #10041 available at: 10041: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf</a>   |
| Regulatory                       | For research purposes only  |

# **Related Products**

**Recommended Negative Controls** 

MOUSE IgG1 NEGATIVE CONTROL:FITC (MCA928F)

## **Recommended Useful Reagents**

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

From March 15, 2021, we will no longer supply printed datasheets with our products. Look out for updates on how to access your digital version at bio-rad-antibodies.com 'M376088:210115'

## Printed on 10 Feb 2021

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint