

## Datasheet: MCA4644

<b>Description:</b>	MOUSE ANTI HUMAN CD155
<b>Specificity:</b>	CD155
<b>Other names:</b>	POLIOVIRUS RECEPTOR
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	SKII.4
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.2 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/25 - 1/200
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunofluorescence	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Human
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> )
<b>Carrier Free</b>	Yes

<b>Approx. Protein Concentrations</b>	IgG concentration 1.0mg/ml
<b>Immunogen</b>	Human SK-N-S1 neuroblastoma cells.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P15151</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">5817</a>    PVR    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	PVS
<b>RRID</b>	AB_1628763
<b>Specificity</b>	<p><b>Mouse anti Human CD155 antibody, clone SKII.4</b> recognizes human CD155, otherwise known as poliovirus receptor (PVR). CD155 is a 397 amino acid type I transmembrane glycoprotein with a predicted molecular mass of ~45 kDa, a member of the nectin family, which acts as a mediator of cell-cell adhesion and cell migration, and is a receptor for poliovirus and cytomegalovirus.</p> <p>CD155 is a primate-restricted marker expressed by monocytes, macrophages, CD34+ thymocytes, endothelial and epithelial cells, and exists as both cell surface (alpha and delta) and secreted (beta and gamma) isoforms. CD155 promotes natural killer (NK) cell adhesion and NK effector functions, interacting with the NK receptors CD96 (Tactile) and CD226 (DNAM-1). CD155 forms a heterodimer with CD113 (nectin-3) which then binds to integrin alphaVbeta3.</p> <p>Mouse anti Human CD155 antibody, clone SKII.4 has been shown to inhibit NK cell-mediated lysis (<a href="#">Fuchs et al. 2004</a>).</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul.
<b>References</b>	<ol style="list-style-type: none"> <li>1. Fuchs, A. <i>et al.</i> (2005) Paradoxical inhibition of human natural interferon-producing cells by the activating receptor NKp44. <a href="#">Blood. 106 (6): 2076-82.</a></li> <li>2. Boles, K.S. <i>et al.</i> (2009) A novel molecular interaction for the adhesion of follicular CD4 T cells to follicular DC. <a href="#">Eur J Immunol. 39 (3): 695-703.</a></li> <li>3. Cella, M. <i>et al.</i> (2010) Loss of DNAM-1 contributes to CD8+ T-cell exhaustion in chronic HIV-1 infection. <a href="#">Eur J Immunol. 40 (4): 949-54.</a></li> </ol>
<b>Storage</b>	<p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.</p>

<b>Guarantee</b>	12 months from date of despatch
<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA4644">https://www.bio-rad-antibodies.com/SDS/MCA4644</a> 10040
<b>Regulatory</b>	For research purposes only

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	<a href="#">RPE</a>
Goat Anti Mouse IgG IgA IgM (STAR87...)	<a href="#">Alk. Phos.</a> , <a href="#">HRP</a>
Goat Anti Mouse IgG (STAR76...)	<a href="#">RPE</a>
Rabbit Anti Mouse IgG (STAR13...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (STAR70...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (H/L) (STAR117...)	<a href="#">Alk. Phos.</a> , <a href="#">DyLight®488</a> , <a href="#">DyLight®550</a> , <a href="#">DyLight®650</a> , <a href="#">DyLight®680</a> , <a href="#">DyLight®800</a> , <a href="#">FITC</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR9...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (Fc) (STAR120...)	<a href="#">FITC</a> , <a href="#">HRP</a>
Goat Anti Mouse IgG (STAR77...)	<a href="#">HRP</a>

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)  
'M384096:210513'

Printed on 19 Jan 2024