

Datasheet: MCA4644F

Description:	MOUSE ANTI HUMAN CD155:FITC
Specificity:	CD155
Other names:	POLIOVIRUS RECEPTOR
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	SKII.4
Isotype:	lgG1

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	-			Neat - 1/10

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human			
Product Form	Purified IgG conjuga	1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nr	m)
	FITC	490	525	
Preparation	Purified IgG prepare supernatant	d by affinity chromatog	raphy on Protein G	G from tissue culture
Buffer Solution	Phosphate buffered	saline		
Preservative	0.09% Sodium Azide	e (NaN ₃)		
Stabilisers	1% Bovine Serum A	lbumin		
Approx. Protein Concentrations	IgG concentration 0.	1mg/ml		

ı	m	m	u	n	o	a	e	n
•	•••	•••	•	••	•	3	•	

Human SK-N-S1 neuroblastoma cells.

External Database

Links

UniProt:

P15151 Related reagents

Entrez Gene:

5817 PVR Related reagents

Synonyms

PVS

RRID

AB 1719984

Specificity

Mouse anti Human CD155 antibody, clone SKII.4 recognizes human CD155, otherwise known as poliovirus receptor (PVR). CD155 is a 397 amino acid type I transmembrane glycoprotein with a predicted molecular mass of ~45 kDa, a member of the nectin family, which acts as a mediator of cell-cell adhesion and cell migration, and is a receptor for poliovirus and cytomegalovirus.

CD155 is a primate-restricted marker expressed by monocytes, macrophages, CD34+ thymocytes, endothelial and epithelial cells, and exists as both cell surface (alpha and delta) and secreted (beta and gamma) isoforms. CD155 promotes natural killer (NK) cell adhesion and NK effector functions, interacting with the NK receptors CD96 (Tactile) and CD226 (DNAM-1). CD155 forms a heterodimer with CD113 (nectin-3) which then binds to integrin alphaVbeta3.

Mouse anti Human CD155 antibody, clone SKII.4 has been shown to inhibit NK cell-mediated lysis (Fuchs et al. 2004).

Flow Cytometry

Use 10ul of the suggested working dilution to label 1x10⁶ cells in 100ul.

References

- 1. Fuchs, A. *et al.* (2005) Paradoxic inhibition of human natural interferon-producing cells by the activating receptor NKp44. <u>Blood. 106 (6): 2076-82.</u>
- 2. Boles, K.S. *et al.* (2009) A novel molecular interaction for the adhesion of follicular CD4 T cells to follicular DC. <u>Eur J Immunol.</u> 39 (3): 695-703.
- 3. Cella, M. *et al.* (2010) Loss of DNAM-1 contributes to CD8+ T-cell exhaustion in chronic HIV-1 infection. <u>Eur J Immunol. 40 (4): 949-54.</u>

Storage

Store at +4°C or at -20°C if preferred.

Storage in frost-free freezers is not recommended.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety

Material Safety Datasheet documentation #10041 available at:

Information 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf

Regulatory For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:FITC (MCA928F)

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South Tel: +1 800 265 7376

Worldwide

Tel: +44 (0)1865 852 700 **Europe**

Tel: +49 (0) 89 8090 95 21

America

Fax: +1 919 878 3751

Fax: +44 (0)1865 852 739

Fax: +49 (0) 89 8090 95 50

Tax. 11010

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

'M367796:200529'

Printed on 04 Jan 2021

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint