

Datasheet: MCA4644FT

Description:	MOUSE ANTI HUMAN CD155:FITC
Specificity:	CD155
Other names:	POLIOVIRUS RECEPTOR
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	SKII.4
lsotype:	lgG1
Quantity:	25 µg

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .							
		Yes	No	Not Determined	Suggested Dilution			
	Flow Cytometry	•						
	Where this product has not been tested for use in a particular technique this does not							
	necessarily exclude its a guide only. It is recor system using appropria	mmended that	the use	r titrates the product f	g dilutions are given as or use in their own			
Target Species	Human							
Product Form	Purified IgG conjugate	d to Fluoresce	ein Isothi	ocyanate Isomer 1 (F	ITC) - liquid			
Max Ex/Em	Fluorophore	Excitation Ma	x (nm)	Emission Max (nm)				
	FITC	490		525				
Preparation	Purified IgG prepared supernatant	by affinity chro	omatogra	aphy on Protein G fror	n tissue culture			
Buffer Solution	Phosphate buffered sa							
Preservative Stabilisers	0.09% Sodium Azide(1% Bovine Serum Albo							
Approx. Protein Concentrations	IgG concentration 0.1r	ng/ml						

Immunogen	Human SK-N-S1 neuroblastoma cells.
External Database Links	UniProt: <u>P15151</u> <u>Related reagents</u> Entrez Gene: <u>5817</u> PVR <u>Related reagents</u>
Synonyms	PVS
RRID	AB_2284665
Specificity	 Mouse anti Human CD155 antibody, clone SKII.4 recognizes human CD155, otherwise known as poliovirus receptor (PVR). CD155 is a 397 amino acid type I transmembrane glycoprotein with a predicted molecular mass of ~45 kDa, a member of the nectin family, which acts as a mediator of cell-cell adhesion and cell migration, and is a receptor for poliovirus and cytomegalovirus. CD155 is a primate-restricted marker expressed by monocytes, macrophages, CD34+ thymocytes, endothelial and epithelial cells, and exists as both cell surface (alpha and delta) and secreted (beta and gamma) isoforms. CD155 promotes natural killer (NK) cell adhesion and NK effector functions, interacting with the NK receptors CD96 (Tactile) and CD226 (DNAM-1). CD155 forms a heterodimer with CD113 (nectin-3) which then binds to integrin alphaVbeta3. Mouse anti Human CD155 antibody, clone SKII.4 has been shown to inhibit NK cell-mediated lysis (Fuchs <i>et al.</i> 2004).
Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.
References	 Fuchs, A. <i>et al.</i> (2005) Paradoxic inhibition of human natural interferon-producing cells by the activating receptor NKp44. <u>Blood. 106 (6): 2076-82.</u> Boles, K.S. <i>et al.</i> (2009) A novel molecular interaction for the adhesion of follicular CD4 T cells to follicular DC. <u>Eur J Immunol. 39 (3): 695-703.</u> Cella, M. <i>et al.</i> (2010) Loss of DNAM-1 contributes to CD8+ T-cell exhaustion in chronic HIV-1 infection. <u>Eur J Immunol. 40 (4): 949-54.</u>
Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety	Material Safety Datasheet documentation #10041 available at:

Informati	on 10041: j	https://www.bio	<u>S/10041.pdf</u>					
Regulato	ry For rese	For research purposes only						
Relate	d Products							
Recomn	nended Negative C	ontrols						
MOUSE Ig	G1 NEGATIVE CONTR	OL:FITC (MCAS	9 <u>28F)</u>					
Recomm	nended Useful Rea	gents						
	EROBLOCK (BUF070A EROBLOCK (BUF070B	-						
North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50			
America	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad.com		Email: antibody_sales_de@bio-rad.com			
			'M367797:200529'					

Printed on 04 Jan 2021

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint