

Datasheet: MCA4659F

Description:	HAMSTER ANTI MOUSE KLRG1:FITC
Specificity:	KLRG1
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	2F1
Isotype:	IgG
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.1mg/ml		
Immunogen	Activated NK (A-LAK) cells from B6 (H-2b) mice.		

External Database**Links****UniProt:**[O88713](#)[Related reagents](#)**Entrez Gene:**[50928](#)

Klrg1

[Related reagents](#)

Synonyms

Mafa

RRID

AB_10598752

Fusion Partners

Popliteal lymph node cells from immunized Syrian hamsters were fused with cells of the P3X63Ag8.653 plasmacytoma cell line.

Specificity

Hamster anti Mouse KLRG1 antibody, clone 2F1 recognizes mouse KLRG1 (killer cell lectin-like receptor G1), the mouse homolog of the rat mast cell function-associated antigen (MAFA). It is expressed by activated CD8 T-cells and by a large proportion of natural killer (NK) cells. It is not found in the spleen, thymus, lymph nodes, testis, brain or kidney and is not expressed on mast cell lines, bone-marrow-derived mast cells or peritoneal mast cells. T-cells expressing KLRG1 have greatly reduced proliferation potential, but are not limited in their capacity to perform immediate effector cell functions, such as the secretion of gamma interferon. KLRG1 plays an inhibitory role on NK cells and T-cells by binding to their ligands.

Flow Cytometry

Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.

The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR ([BUF041A/B](#)).

References

1. Hanke, T. *et al.* (1998) 2F1 antigen, the mouse homolog of the rat "mast cell function-associated antigen", is a lectin-like type II transmembrane receptor expressed by natural killer cells. [Eur J Immunol. 28 \(12\): 4409-17.](#)
2. Rosshart, S. *et al.* (2008) Interaction of KLRG1 with E-cadherin: new functional and structural insights. [Eur J Immunol. 38 \(12\): 3354-64.](#)
3. Beyersdorf, N. *et al.* (2007) Characterization of mouse CD4 T cell subsets defined by expression of KLRG1. [Eur J Immunol. 37 \(12\): 3445-54.](#)

Storage

Store at +4°C or at -20°C if preferred.

Storage in frost-free freezers is not recommended.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Acknowledgements

This product may be used for research purposes only. It is not licensed for resale and may only be used by the buyer. This product may not be used and is not licensed for clinical

assays, where the results of such assays are provided as a diagnostic service. If a diagnostic or therapeutic use is anticipated, then a license must be requested from the University of California. The availability of such diagnostic and therapeutic use license(s) cannot be guaranteed from the University of California.

Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Negative Controls

[HAMSTER \(ARMENIAN\) IgG NEGATIVE CONTROL:FITC \(MCA2356F\)](#)

Recommended Useful Reagents

[MOUSE SEROBLOCK FcR \(BUF041A\)](#)

[MOUSE SEROBLOCK FcR \(BUF041B\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

From March 15, 2021, we will no longer supply printed datasheets with our products.
Look out for updates on how to access your digital version at [bio-rad-antibodies.com](https://www.bio-rad-antibodies.com)

'M373536:200923'

Printed on 10 Feb 2021

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)