

Datasheet: MCA465GA

BATCH NUMBER 171068

Description:	MOUSE ANTI HUMAN VINCULIN
Specificity:	VINCULIN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	V284
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	▪			
Immunohistology - Frozen	▪			
Immunohistology - Paraffin (2)	▪			
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting	▪			1/500 - 1/2000
Immunofluorescence	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) Membrane permeabilization is required for this application. The use of Leucoperm (Product Code [BUF09](#)) is recommended for this purpose.

(2) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.

Target Species

Human

Species Cross Reactivity

Reacts with: Rat, Rabbit, Chicken, Mouse

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for

further information.

Product Form	Purified IgG - liquid
---------------------	-----------------------

Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
--------------------	---

Buffer Solution	Phosphate buffered saline
------------------------	---------------------------

Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
---------------------------------	--

Carrier Free	Yes
---------------------	-----

Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
---------------------------------------	-----------------------------

Immunogen	Purified vinculin from human platelets.
------------------	---

External Database Links	UniProt: P18206 Related reagents Entrez Gene: 7414 VCL Related reagents
--------------------------------	--

Fusion Partners	Spleen cells of immunized mice were fused with cells from the SP2/0 mouse myeloma cell line.
------------------------	--

Specificity	<p>Mouse anti Human vinculin antibody, clone V284 recognizes human vinculin, also known as metavinculin. Vinculin is a highly conserved, cytoskeletal protein associated with cell-cell and cell-matrix adhesion. Vinculin is an actin filament (F-actin)-binding protein involved in anchoring F-actin to the membrane.</p> <p>Mouse anti Human vinculin antibody, clone V284 detects vinculin as a ~130 kDa band in homogenates of platelets and other cells. It reacts with leucocytes, muscle cells, epithelial cells and fibroblasts.</p>
--------------------	---

Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul
-----------------------	--

Histology Positive Control Tissue	Human appendix
--	----------------

References	<ol style="list-style-type: none">Porter, R.M. <i>et al.</i> (1993) Monoclonal antibodies to cytoskeletal proteins: an immunohistochemical investigation of human colon cancer. J Pathol. 170 (4): 435-40.Kubler, M.D. & Watt, F.M. (1993) Changes in the distribution of actin-associated proteins during epidermal wound healing. J Invest Dermatol. 100 (6): 785-9.DeFife, K.M. <i>et al.</i> (1999) Cytoskeletal and adhesive structural polarizations accompany IL-13-induced human macrophage fusion. J Histochem Cytochem. 47: 65-74.
-------------------	--

4. Jester, J.V. *et al.* (1999) Transforming growth factor(beta)-mediated corneal myofibroblast differentiation requires actin and fibronectin assembly. [Invest Ophthalmol Vis Sci. 40: 1959-67.](#)
5. Dlugosz, J.A. *et al.* (2000) Stretch-induced mesangial cell ERK1/ERK2 activation is enhanced in high glucose by decreased dephosphorylation. [Am J Physiol Renal Physiol. 279: F688-97.](#)
6. Brown, H. *et al.* (2001) Blood-brain barrier function in cerebral malaria in Malawian children. [Am J Trop Med Hyg. 64 \(3-4\): 207-13.](#)
7. Frangogiannis, N.G. *et al.* (2002) Active interstitial remodeling: an important process in the hibernating human myocardium. [J Am Coll Cardiol. 39: 1468-74.](#)
8. Jester, J.V. *et al.* (2003) Myofibroblast differentiation of normal human keratocytes and hTERT, extended-life human corneal fibroblasts. [Invest Ophthalmol Vis Sci. 44: 1850-8.](#)
9. Hua, H. *et al.* (2003) Endothelin-1 activates mesangial cell ERK1/2 via EGF-receptor transactivation and caveolin-1 interaction. [Am J Physiol Renal Physiol. 284: F303-12.](#)
10. Frye, M. *et al.* (2003) Evidence that Myc activation depletes the epidermal stem cell compartment by modulating adhesive interactions with the local microenvironment. [Development. 130: 2793-808.](#)
11. Monsees, T.K. *et al.* (2005) Effects of different titanium alloys and nanosize surface patterning on adhesion, differentiation, and orientation of osteoblast-like cells. [Cells Tissues Organs. 180: 81-95.](#)
12. Rebhun, R.B. *et al.* (2006) Targeting receptor tyrosine kinase on lymphatic endothelial cells for the therapy of colon cancer lymph node metastasis. [Neoplasia. 8: 747-57.](#)
13. Zheng, P.P. *et al.* (2007) Hela I-CaD undergoes a DNA replication-associated switch in localization from the cytoplasm to the nuclei of endothelial cells/endothelial progenitor cells in human tumor vasculature. [Cancer Biol Ther. 6: 886-90.](#)
14. Zheng, P.P. *et al.* (2007) Hela I-CaD is implicated in the migration of endothelial cells/endothelial progenitor cells in human neoplasms. [Cell Adh Migr. 1: 84-91.](#)
15. Gray, M.J. *et al.* (2008) Therapeutic targeting of Id2 reduces growth of human colorectal carcinoma in the murine liver. [Oncogene. 27: 7192-200.](#)
16. Andrews, K.D. *et al.* (2008) Vascular prostheses: performance related to cell-shear responses. [J Surg Res. 149: 39-46.](#)
17. Rio, M. *et al.* (2016) Microfluidic measurement of cell motility in response to applied non-homogeneous DC electric fields [Journal of Sensors and Sensor Systems. 5 \(2\): 237-43.](#)
18. Ananthaseshan, S. *et al.* (2017) Locally Transplanted CD34+ Bone Marrow-Derived Cells Contribute to Vascular Healing After Vascular Injury. [Transplant Proc. 49 \(6\): 1467-76.](#)
19. May, C.A. (2022) Aponeurosis linguae-Myocutaneous or myotendinous junctions of skeletal muscle fibres in the human tongue? [J Anat. 241 \(1\): 168-172.](#)
20. Bola, S. *et al.* (2023) Analysis of Electric Field Stimulation in Blue Light Stressed 661W Cells [Int J Mol Sci. 24 \(4\): 3433.](#)
21. Ran, X. *et al.* (2023) Modulation of the hippo-YAP pathway by cyclic stretch in rat type 2 alveolar epithelial cells-a proof-of-concept study. [Front Physiol. 14: 1253810.](#)
22. Simiczjew, A. *et al.* (2025) Phenotype switching in highly invasive resistant to vemurafenib and cobimetinib melanoma cells. [Cell Commun Signal. 23 \(1\): 449.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at

-20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
------------------	---------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA465GA
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87...)	HRP
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Rabbit Anti Mouse IgG (STAR12...)	RPE
Rabbit Anti Mouse IgG (STAR13...)	HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M437706:250318'

Printed on 26 May 2026