

Datasheet: MCA4660F

Description:	HAMSTER ANTI MOUSE CD137:FITC
Specificity:	CD137
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	17B5
Isotype:	lgG
Quantity:	0.1 mg

Entrez Gene:

21942 Tnfrsf9

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				Neat - 1/10

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse					
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid					
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)			
	FITC	490	525			
Preparation	Purified IgG prep	ared by affinity chromatog	raphy on Protein G fror	m tissue culture supernata		
Buffer Solution	Phosphate buffer	ed saline				
Preservative	0.09% Sodium Azide (NaN ₃)					
Stabilisers	1% Bovine Serun	n Albumin				
Approx. Protein Concentrations	IgG concentration	n 0.1mg/ml				
External Database Links	UniProt: <u>P20334</u> R	elated reagents				

Related reagents

Synonyms	Cd137, Ila, Ly63
RRID	AB_10548157
Specificity	Hamster anti Mouse CD137 antibody, clone 17B5 recognizes mouse CD137, also known as 4-1BB ligand receptor, a member of the tumour necrosis factor superfamily. CD137 is a potent costimulatory receptor for the growth and functional maturation of activated T-cells in the presence of T-cell receptor signal. It is found on activated T-cells, NK cells, monocytes and dendritic cells. Signalling through CD137 triggers T-cell expansion, cytokine production and prevention of activation-induced death of effector T-cells. CD137-mediated activation is distinct from and independent of CD28. Antibody cross-linking of CD137 on NK cells stimulates the secretion of IFN-gamma, cellular proliferation, CD25 expression and responsiveness to IL-2 and is associated with tumour activity, but not cytotoxicity.
	Hamster anti Mouse CD137 antibody, clone 17B5 may be used to block CD137-mediated T-cell proliferation.
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.
	The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR (<u>BUF041A/B</u>).
References	1. Zheng, G. <i>et al.</i> (2004) The 4-1BB costimulation augments the proliferation of CD4+CD25+ regulatory T cells. J Immunol. 173 (4): 2428-34.
Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Acknowledgements	This product may be used for research purposes only. It is not licensed for resale and may only be used by the buyer. This product may not be used and is not licensed for clinical assays, where the results of such assays are provided as a diagnostic service. If a diagnostic or therapeutic use is anticipated, then a license must be requested from the University of California. The availability of such diagnostic and therapeutic use license(s) cannot be guaranteed from the University of California.
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

HAMSTER (ARMENIAN) IgG NEGATIVE CONTROL:FITC (MCA2356F)

North & South Tel: +1 800 265 7376 America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 'M367830:200529'

Printed on 11 Aug 2020

© 2020 Bio-Rad Laboratories Inc | Legal | Imprint