

## Datasheet: MCA4660GA

<b>Description:</b>	HAMSTER ANTI MOUSE CD137
<b>Specificity:</b>	CD137
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	17B5
<b>Isotype:</b>	IgG
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Mouse
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> )
<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0mg/ml
<b>External Database Links</b>	<b>UniProt:</b>

**Entrez Gene:**

[21942](#)   Tnfrsf9   [Related reagents](#)

---

**Synonyms**      Cd137, Ila, Ly63

---

**RRID**            AB\_2240760

---

**Specificity**      **Hamster anti Mouse CD137 antibody, clone 17B5** recognizes mouse CD137, also known as 4-1BB ligand receptor, a member of the tumour necrosis factor superfamily. CD137 is a potent costimulatory receptor for the growth and functional maturation of activated T-cells in the presence of T-cell receptor signal. It is found on activated T-cells, NK cells, monocytes and dendritic cells. Signalling through CD137 triggers T-cell expansion, cytokine production and prevention of activation-induced death of effector T-cells. CD137-mediated activation is distinct from and independent of CD28. Antibody cross-linking of CD137 on NK cells stimulates the secretion of IFN-gamma, cellular proliferation, CD25 expression and responsiveness to IL-2 and is associated with tumour activity, but not cytotoxicity.

Hamster anti Mouse CD137 antibody, clone 17B5 may be used to block CD137-mediated T-cell proliferation.

---

**Flow Cytometry**      Use 10ul of the suggested working dilution to label  $1 \times 10^6$  cells in 100ul.

---

**References**            1. Zheng, G. *et al.* (2004) The 4-1BB costimulation augments the proliferation of CD4+CD25+ regulatory T cells. [J Immunol. 173 \(4\): 2428-34.](#)

---

**Storage**              Store at +4°C or at -20°C if preferred.  
Storage in frost-free freezers is not recommended.  
This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee**            12 months from date of despatch

---

**Acknowledgements**      This product may be used for research purposes only. It is not licensed for resale and may only be used by the buyer. This product may not be used and is not licensed for clinical assays, where the results of such assays are provided as a diagnostic service. If a diagnostic or therapeutic use is anticipated, then a license must be requested from the University of California. The availability of such diagnostic and therapeutic use license(s) cannot be guaranteed from the University of California.

---

**Health And Safety Information**      Material Safety Datasheet documentation #10040 available at:  
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

---

**Regulatory**            For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Goat Anti Hamster IgG (STAR104...) [DyLight@800](#), [FITC](#)

Goat Anti Hamster IgG (STAR79...) [Biotin](#), [FITC](#), [HRP](#)

### Recommended Negative Controls

[HAMSTER \(ARMENIAN\) IgG NEGATIVE CONTROL \(MCA2356\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

From March 15, 2021, we will no longer supply printed datasheets with our products.  
Look out for updates on how to access your digital version at [bio-rad-antibodies.com](http://bio-rad-antibodies.com)

'M367831:200529'

**Printed on 12 Feb 2021**

---

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)