

Datasheet: MCA4662 BATCH NUMBER 168961

| Description: | MOUSE ANTI HUMAN DECTIN-1 | | |
|---------------|-----------------------------|--|--|
| Specificity: | DECTIN-1 | | |
| Other names: | BETA GLUCAN RECEPTOR, CD369 | | |
| Format: | Purified | | |
| Product Type: | Monoclonal Antibody | | |
| Clone: | BD6 | | |
| Isotype: | lgG2b | | |
| Quantity: | 0.2 mg | | |
| | | | |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------------------|-----|----|----------------|--------------------|
| Flow Cytometry | • | | | 1/25 - 1/200 |
| Immunohistology - Frozen | | | | |
| Immunohistology - Paraffin | | | | |
| ELISA | | | | |
| Immunoprecipitation | | | | |
| Western Blotting | | | • | |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

| Target Species | Human | |
|-----------------------------|---|-----------------------|
| Product Form | Purified IgG - liquid | |
| Preparation | Purified IgG prepared by affinity chromatography on Protein A supernatant | A from tissue culture |
| Buffer Solution | Phosphate buffered saline | |
| Preservative Stabilisers | 0.09% Sodium Azide (NaN ₃) | |

| Carrier Free | Yes |
|-----------------------------------|--|
| Approx. Protein Concentrations | IgG concentration 1.0mg/ml |
| Immunogen | NIH3T3 cells expressing full length Beta-glucan receptor. |
| External Database Links | UniProt: Q9BXN2 Related reagents Entrez Gene: 64581 CLEC7A Related reagents |
| Synonyms | BGR, CLECSF12, DECTIN1 |
| RRID | AB_1628802 |
| Fusion Partners | Spleen cells from immunized C57BL/6 were fused with cells of the murine SP2 myeloma cell line. |
| Specificity | Mouse anti Human Dectin-1 antibody, clone BD6 recognizes human Dectin-1, also known as beta-glucan receptor (β GR) or CD369. Dectin-1 is a C-type lectin domain family member. It is a major pathogen pattern-recognition receptor that binds beta-glucans, glucose polymers that form cell-wall components or exopolymers of yeasts, fungi and some bacteria. Beta-glucans have also been used experimentally and therapeutically as immunomodulators that enhance resistance to bacterial, yeast, viral and protozoan infections, as well as tumor formation. Dectin-1 also acts as a co-stimulatory molecule on T-cells to induce their proliferation and is necessary for the TLR2-mediated inflammatory response. It is highly expressed on peripheral blood leukocytes and dendritic cells. Human Dectin-1 has two major (β GR-A and β GR-B) and several minor isoforms. β GR-A and β GR-B differs by the presence and absence, respectively, of a stalk region. Mouse anti Human Dectin-1 antibody, clone BD6 recognizes β GR-A only. |
| Flow Cytometry | Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul. |
| References | Plantinga, T.S. <i>et al.</i> (2009) Early stop polymorphism in human DECTIN-1 is associated with increased candida colonization in hematopoietic stem cell transplant recipients. <u>Clin Infect Dis. 49 (5): 724-32.</u> Hoeksema, M.A. <i>et al.</i> (2014) Targeting macrophage Histone deacetylase 3 stabilizes atherosclerotic lesions. <u>EMBO Mol Med. 6 (9): 1124-32.</u> |
| Storage | This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. |
| | Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. |

| Guarantee | 12 months from date of despatch |
|----------------------------------|--|
| Health And Safety Information | Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA4662 10040 |
| Regulatory | For research purposes only |

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)

Goat Anti Mouse IgG IgA IgM (STAR87...)

RPE

Goat Anti Mouse IgG (STAR76...)

RPE

Goat Anti Mouse IgG (STAR70...)

FITC

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Goat Anti Mouse IgG (STAR77...) <u>HRP</u>

Rabbit Anti Mouse IgG (STAR9...) <u>FITC</u>

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

Rabbit Anti Mouse IgG (STAR13...) HRP

Recommended Negative Controls

MOUSE IgG2b NEGATIVE CONTROL (MCA691)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

 $\textbf{Email: antibody_sales_us@bio-rad.com} \\ \textbf{Email: antibody_sales_uk@bio-rad.com} \\ \textbf{Email: antibody_sales_uk@b$

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M431479:240717'

Printed on 17 Jul 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint