

Datasheet: MCA4677

BATCH NUMBER 1804

Description:	MOUSE ANTI HUMAN FACTOR VIII
Specificity:	FACTOR VIII
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	RFF-VIII C/8
Isotype:	IgG1
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			1/500 - 1/15000
Immunoprecipitation			▪	
Western Blotting	▪			
Radioimmunoassays	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Species Cross Reactivity	<p>Reacts with: Pig</p> <p>Does not react with: Mouse, Dog, Rat</p> <p>N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
Product Form	Purified IgG - liquid

Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
Immunogen	Affinity purified human Factor VIII.
External Database Links	<p>UniProt: P00451 Related reagents</p> <p>Entrez Gene: 2157 F8 Related reagents</p>
Synonyms	F8C
RRID	AB_1833762
Fusion Partners	Spleen cells from an immunized Balb/c mouse we fused both cells of the P3-NS/a-Ag4-1 mouse myeloma.
Specificity	<p>Mouse anti Human Factor VIII antibody, clone RFF-VIII C/8 recognizes human Factor VIII, an essential blood coagulation factor. Whilst circulating in the blood, it is mostly stably complexed to von Willebrand factor. It is activated through cleavage at various sites, dissociates from the complex and interacts with Factor IXa, in the presence of calcium ions and phospholipids, to convert Factor X to the activated Factor Xa, which activates thrombin. Thrombin cleaves fibrinogen into fibrin, which polymerizes and cross-links to form a blood clot. The activated Factor VIII is proteolytically inactivated and cleared from the bloodstream.</p> <p>Defects in Factor VIII cause haemophilia A, a disorder characterized by the body's inability to control blood clotting. This could result in severe blood loss, even with minor injuries.</p> <p>Mouse anti Human Factor VIII antibody, clone RFF-VIII C/8 is a very potent coagulation inhibitor. It recognizes an epitope towards the N-terminus of full-length Factor VIII. It also recognizes the 210 kDa, 90 kDa and 40 kDa cleavage products. Mouse anti Human Factor VIII antibody, clone RFF-VIII C/8 does not cross-react with von Willebrand factor.</p>
References	<ol style="list-style-type: none"> 1. Rotblat, F. <i>et al.</i> (1983) Monoclonal antibodies to human procoagulant factor VIII. J Lab Clin Med. 101 (5): 736-46. 2. Tiarks, C. <i>et al.</i> (1987) Identification of six functional clotting factor VIII:C epitopes by

analysis of cross-reactive public idiotypes in murine monoclonal VIII:C inhibitors. [Thromb Res. 45 \(5\): 527-37.](#)

3. Purohit, V.S. *et al.* (2006) Influence of aggregation on immunogenicity of recombinant human Factor VIII in hemophilia A mice. [J Pharm Sci. 95 \(2\): 358-71.](#)

4. Zaniboni, A. *et al.* (2015) In vitro differentiation of porcine aortic vascular precursor cells to endothelial and vascular smooth muscle cells. [Am J Physiol Cell Physiol. 309 \(5\): C320-31.](#)

5. Turner NA & Moake JL (2015) Factor VIII Is Synthesized in Human Endothelial Cells, Packaged in Weibel-Palade Bodies and Secreted Bound to ULVWF Strings. [PLoS One. 10 \(10\): e0140740.](#)

Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA4677 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (STAR70...)	FITC
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M367865:200529'

Printed on 13 May 2025

