

## Datasheet: MCA4678G BATCH NUMBER 152379

| Description:  | MOUSE ANTI HUMAN FACTOR VIII |
|---------------|------------------------------|
| Specificity:  | FACTOR VIII                  |
| Format:       | Purified                     |
| Product Type: | Monoclonal Antibody          |
| Clone:        | RFF-VIIIC/5                  |
| lsotype:      | lgG1                         |
| Quantity:     | 0.2 mg                       |

## **Product Details**

| Applications                | This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> . |                   |    |                |                    |  |
|-----------------------------|--|-------------------|----|----------------|--------------------|--|
|                             |  | Yes               | No | Not Determined | Suggested Dilution |  |
|                             | Flow Cytometry   |                   |    |                |                    |  |
|                             | Immunohistology - Frozen   |                   |    |                |                    |  |
|                             | Immunohistology - Paraffin   |                   |    | •              |                    |  |
|                             | ELISA  | -                 |    |                | 1/400 - 1/4000     |  |
|                             | Immunoprecipitation  |                   |    | •              |                    |  |
|                             | Western Blotting   | -                 |    |                |                    |  |
|                             | Radioimmunoassays  | -                 |    |                |                    |  |
|                             | necessarily exclude its use in such procedures. Suggested working dilutions ar<br>a guide only. It is recommended that the user titrates the product for use in the<br>system using appropriate negative/positive controls.  |                   |    |                |                    |  |
| Target Species              | Human  |                   |    |                |                    |  |
| Product Form                | Purified IgG - liquid  |                   |    |                |                    |  |
| Preparation                 | Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant  |                   |    |                |                    |  |
| Buffer Solution             | Phosphate buffered salin   | e                 |    |                |                    |  |
| Preservative<br>Stabilisers | 0.09% Sodium Azide (Na   | ıN <sub>3</sub> ) |    |                |                    |  |

| Carrier Free                      | Yes  |  |  |  |
|-----------------------------------|--|--|--|--|
| Approx. Protein<br>Concentrations | IgG concentration 1.0mg/ml   |  |  |  |
| Immunogen                         | Affinity purified human Factor VIII.   |  |  |  |
| External Database<br>Links        | UniProt:         P00451       Related reagents         Entrez Gene:         2157       F8       Related reagents   |  |  |  |
| Synonyms                          | F8C  |  |  |  |
| RRID                              | AB_1833761   |  |  |  |
| Fusion Partners                   | Spleen cells from immunized Balb/c mice were fused with P3-NS1/1-Ag4-1 myeloma cells.  |  |  |  |
| Specificity                       | <ul> <li>Mouse anti Human Factor VIII antibody, clone RFF-VIIIC/5 recognizes human Factor VIII, an essential blood coagulation factor. Factor VIII is primarily synthesized in the liver and is secreted into the bloodstream where it forms a complex with von Willebrand factor. It is activated through cleavage at various sites, dissociates from the complex and interacts with Factor IXa, in the presence of calcium ions and phospholipids, to convert Factor X to the activated Factor Xa, which in turn activates thrombin. Thrombin cleaves fibrinogen into fibrin, which polymerises and cross-links to form a blood clot. The activated Factor VIII is proteolytically inactivated and cleared from the bloodstream.</li> <li>Defects in Factor VIII cause haemophilia A (HEMA), a disorder characterised by the body's inability to control blood clotting, potentially resulting in severe blood loss, even with minor injuries.</li> <li>Mouse anti Human Factor VIII antibody, clone RFF-VIIIC/5 recognizes an epitope towards the C-terminus of full-length Factor VIII. It also recognizes the ~80 kDa and ~70 kDa cleavage products.</li> </ul> |  |  |  |
| References                        | <ol> <li>Rotblat, F. <i>et al.</i> (1983) Monoclonal antibodies to human procoagulant factor VIII. <u>J Lab</u><br/><u>Clin Med. 101 (5): 736-46.</u></li> <li>Tiarks, C. <i>et al.</i> (1987) Identification of six functional clotting factor VIII:C epitopes by<br/>analysis of cross-reactive public idiotypes in murine monoclonal VIII:C inhibitors. <u>Thromb</u><br/><u>Res. 45 (5): 527-37.</u></li> </ol>  |  |  |  |
| Storage                           | Store at +4°C or at -20°C if preferred.<br>Storage in frost-free freezers is not recommended.<br>This product should be stored undiluted. Avoid repeated freezing and thawing as this may<br>denature the antibody. Should this product contain a precipitate we recommend<br>microcentrifugation before use.  |  |  |  |

| 12 months from date of despatch   |  |  |  |
|---|--|--|--|
| Material Safety Datasheet documentation #10040 available at <a href="https://www.bio-rad-antibodies.com/SDS/MCA4678G">https://www.bio-rad-antibodies.com/SDS/MCA4678G</a> 10040 |  |  |  |
| For research purposes only  |  |  |  |
| Related Products  |  |  |  |
| econdary Antibodies   |  |  |  |
|   |  |  |  |

| Rabbit Anti Mouse IgG (STAR12)   | RPE  |  |  |  |  |
|--|--|--|--|--|--|
| Goat Anti Mouse IgG IgA IgM (STAR87  | .) <u>HRP</u>  |  |  |  |  |
| Goat Anti Mouse IgG (STAR76)   | RPE  |  |  |  |  |
| Goat Anti Mouse IgG (STAR70)   | <u>FITC</u>  |  |  |  |  |
| Rabbit Anti Mouse IgG (STAR13)   | HRP  |  |  |  |  |
| Goat Anti Mouse IgG (Fc) (STAR120)   | FITC, HRP  |  |  |  |  |
| Rabbit Anti Mouse IgG (STAR9)  | <u>FITC</u>  |  |  |  |  |
| Goat Anti Mouse IgG (STAR77)   | HRP  |  |  |  |  |
| Goat Anti Mouse IgG (H/L) (STAR117)  | Alk. Phos., DyLight®488, DyLight®550,  |  |  |  |  |
|  | DyLight®650, DyLight®680, DyLight®800,   |  |  |  |  |
|  | FITC, HRP  |  |  |  |  |
|  |  |  |  |  |  |
| North & South         Tel: +1 800 265 7376         Worldwin           America         Fax: +1 919 878 3751 | ide         Tel: +44 (0)1865 852 700         Europe         Tel: +49 (0) 89 8090 95 21           Fax: +44 (0)1865 852 739         Fax: +49 (0) 89 8090 95 50 |  |  |  |  |

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M367868:200529'

Email: antibody\_sales\_uk@bio-rad.com

Email: antibody\_sales\_de@bio-rad.com

## Printed on 08 May 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint

Email: antibody\_sales\_us@bio-rad.com