

Datasheet: MCA467GA BATCH NUMBER 0713

Description:	MOUSE ANTI HUMAN CD41
Specificity:	CD41
Other names:	INTEGRIN ALPHA IIB
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	PM6/248
Isotype:	lgG1
Quantity:	0.1 mg

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .				
		Yes	No	Not Determined	Suggested Dilution
	Flow Cytometry	-			
	Immunohistology - Frozen				
	Immunohistology - Paraffin				

Western Blotting

Immunoprecipitation

ELISA

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

•

.

•

Target Species	Human
Species Cross Reactivity	Reacts with: Baboon, Mink, Mustelid N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture

	supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	0.09% Sodium Azide		
Carrier Free	Yes		
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml		
Immunogen	Human platelet plasma membranes.		
External Database Links	UniProt: <u>P08514</u> <u>Related reagents</u> Entrez Gene: <u>3674</u> ITGA2B <u>Related reagents</u>		
Synonyms	GP2B, ITGAB		
RRID	AB_324706		
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cel myeloma cell line.	lls of the mouse SP2/0	
Specificity	 Mouse anti Human CD41 antibody, clone PM6/248 recognizes the human CD41 cell surface antigen, a ~140 kDa glycoprotein expressed by platelets and megakaryocytes. CD41 is also known as platelet glycoprotein IIb, and functions as a receptor for fibrinogen, fibronectin and vWF. It has not been established if clone PM6/248 recognizes free CD41 or CD41 only when complexed with CD61. However, antibody binding is reduced in the presence of EDTA suggesting that the epitope recognized is dependent upon an intact CD41/61 complex. 		
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells or	100ul whole blood	
References	 Hornby, E.J. <i>et al.</i> (1991) Activation of human platelets by e antibody, PM6/248, to glycoprotein IIb-IIIa. <u>Br J Haematol. 79 (</u>2. Michelson, A.D. <i>et al.</i> (1995) A panel of platelet mAb for the thrombosis in baboons. Leucocyte Typing V. Oxford University 3. Maloney, S.F. <i>ety al.</i> (2010) P2Y12 or P2Y1 inhibitors reduce microfluidic model of thrombosis while apyrase lacks efficacy u <u>Biol (Camb). 2: 183-92.</u> Aasted, B. <i>et al.</i> (2007) Reactivity of monoclonal antibodies cells from mink. <u>Vet Immunol Immunopathol. 119: 27-37.</u> Massoudy, P. <i>et al.</i> (2001) Evidence for inflammatory respondent. 	(2): 277-85. study of haemostasis and Press p 1230-1231. e platelet deposition in a under flow conditions. <u>Integr</u> to human CD antigens with	

	 coronary artery bypass grafting with cardiopulmonary bypass. <u>Chest. 119: 31-6.</u> 6. Kahng, J. <i>et al.</i> (2008) Quantitative comparisons of antibody-binding sites of platelet glycoprotein IIb/IIIa in aplastic anemia and idiopathic thrombocytopenic purpura. <u>Ann Clin Lab Sci. 38: 6-11.</u> 7. Chae, H. <i>et al.</i> (2009) EDTA Inhibits the Binding of Clone 96.2C1, an Anti-CD41a Monoclonal Antibody, to the Platelets and Addition of Heparin and CaCl2 to the Antibody Neutralizes the EDTA-induced Inhibitory Effect Korean J Hematol 44: 42 - 6.
	 Zahler, S. et al. (1999) Acute cardiac inflammatory responses to postischemic reperfusion during cardiopulmonary bypass. <u>Cardiovasc Res. 41: 722-30.</u>
	9. Wang, J.S. <i>et al.</i> (2005) Effects of exercise training and deconditioning on platelet
	aggregation induced by alternating shear stress in men. <u>Arterioscler Thromb Vasc Biol.</u> 25: 454-60.
	10. Welsh, J.D. <i>et al.</i> (2012) Platelet-targeting sensor reveals thrombin gradients within blood clots forming in microfluidic assays and in mouse. <u>J Thromb Haemost. 10 (11):</u>
	<u>2344-53.</u>
	11. Kamat, V. <i>et al.</i> (2015) Microfluidic assessment of functional culture-derived platelets in human thrombi under flow. <u>Exp Hematol. 43 (10): 891-900.e4.</u>
	12. Vučetić, D. <i>et al.</i> (2018) Flow cytometry analysis of platelet populations: usefulness for monitoringthe storage lesion in pooled buffy-coat platelet concentrates. <u>Blood Transfus. 16</u> (<u>1): 83-92.</u>
	 13. Foruzanmehr, M. <i>et al.</i> (2014) Nano-structure TiO₂ film coating on 316L stainless steel via sol-gel technique for blood compatibility improvement Nanomedicine J. 1 (3): 128-36. 14. Miyazaki, K. <i>et al.</i> (2015) Immature platelet fraction measurement is influenced by platelet size and is a useful parameter for discrimination of macrothrombocytopenia. <u>Hematology. 20 (10): 587-92.</u>
Storage	Store at +4°C or at -20°C if preferred.
	This product should be stored undiluted.
	Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA467GA 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)RPEGoat Anti Mouse IgG IgA IgM (STAR87...)HRP

Goat Anti Mouse IgG (STAR76)	RPE
Goat Anti Mouse IgG (STAR70)	<u>FITC</u>
Rabbit Anti Mouse IgG (STAR13)	HRP
Goat Anti Mouse IgG (Fc) (STAR120)	FITC, HRP
Rabbit Anti Mouse IgG (STAR9)	<u>FITC</u>
Goat Anti Mouse IgG (STAR77)	HRP
Goat Anti Mouse IgG (H/L) (STAR117)	Alk. Phos., DyLight®488, DyLight®550,
	<u>DyLight®650</u> , <u>DyLight®680</u> , <u>DyLight®800</u> ,
	FITC, HRP
Recommended Negative Controls	

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-	rad.com	Email: antibody_sales_uk@bio-	rad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M367875:200529'

Printed on 23 May 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint