

# Datasheet: MCA4685

### **BATCH NUMBER 1612**

Description:	MOUSE ANTI HUMAN CYTOCHROME B245 HEAVY CHAIN		
Specificity:	CYTOCHROME B245 HEAVY CHAIN		
Other names:	gp91phox		
Format:	Purified		
<b>Product Type:</b>	Monoclonal Antibody		
Clone:	NL7		
Isotype:	IgG1		
Quantity:	0.2 mg		

## **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	•			
Immunohistology - Frozen	•			
Immunohistology - Paraffin				
ELISA	•			
Immunoprecipitation				
Western Blotting	•			
Immunofluorescence	•			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1)Membrane permeabilisation is required for this application. Bio-Rad recommends the use of Leucoperm<sup>™</sup> (Product Code <u>BUF09</u>) for this purpose.

Target Species	Human	
Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared by affinity chromatography on Protein C supernatant	3 from tissue culture
Buffer Solution	Phosphate buffered saline	

**Preservative** 0.09% Sodium Azide (NaN<sub>3</sub>) **Stabilisers Carrier Free** Yes Approx. Protein IgG concentration 1.0mg/ml Concentrations Immunogen Cytochrome B245 solubilised in octyl-beta-glucopyranoside. **External Database UniProt:** Links P04839 Related reagents **Entrez Gene:** 1536 CYBB Related reagents **Synonyms** NOX2 **RRID** AB 1963543 **Fusion Partners** Spleen cells from immunised BALB/c were fused with cells of the 3PU1 myeloma cell line. Specificity

Mouse anti Human Cytochrome B245 Heavy Chain antibody, clone NL7 recognizes

the heavy chain of cytochrome B245, also known as gp91phox or NOX2, a subunit of the nicotinamide adenine dinucleotide phosphate-oxidase (NADPH oxidase). NADPH oxidase is a membrane-bound enzyme complex in neutrophils that produces the superoxide anion (O2<sup>-</sup>) by transferring single electrons from NADPH inside the cell across the membrane and coupling them to molecular oxygen. Superoxide is used for the destruction of pathogens. Individuals with defects in NADPH oxidase suffer from chronic granulomatous disease, characterised by severe recurrent infections.

The heavy chain (gp91phox), along with the light chain (p22phox) form cytochrome B245 (also known as flavocytochome b, or Cyt b), which is always membrane-resident. Upon the appropriate stimulus, other components of NADPH oxidase (p47phox, p40phox, p67phox and Rac2) rapidly translocate from the cytosol to the cell membrane to form the complete enzyme complex. Cytochrome B245 appears to serve as a scaffold for the assembly of the other cytosolic units of the oxidase and to provide a transmembrane pathway for electrons.

Mouse anti Human Cytochrome B245 Heavy Chain antibody, clone NL7 binds amino acids 498-506 (EKDVITGLK), a cytosolic epitope on gp91phox not accessible on intact neutrophils. It recognizes the native as well as the denatured form of the protein. Clone NL7 has an inhibitory effect on NADPH oxidase activation. Research indicates that this inhibitory effect is exerted during the initial stages of oxidase activation. The antibody does not interfere with the translocation of the cytosolic subunits of NADPH oxidase to the cell membrane.

#### Flow Cytometry

Use 10ul of the suggested working dilution to label 1x10<sup>6</sup> cells in 100ul.

Western Blotting	MCA4685 detects a broad band of approximately 90kDa
References	1. Burritt, J.B. <i>et al.</i> (2003) Functional epitope on human neutrophil flavocytochrome b558. J Immunol. 170 (12): 6082-9.
	2. Taylor, R.M. <i>et al.</i> (2004) Site-specific inhibitors of NADPH oxidase activity and
	structural probes of flavocytochrome b: characterization of six monoclonal antibodies to the p22phox subunit. <u>J Immunol</u> . 173 (12): 7349-57.
	3. Taylor, R.M. <i>et al.</i> (2007) Cloning, sequence analysis and confirmation of derived gene sequences for three epitope-mapped monoclonal antibodies against human phagocyte flavocytochrome b. Mol Immunol. 44 (4): 625-37.
	4. Cao, S. (2010) Fc gamma receptors are required for NF-kappaB signaling, microglial activation and dopaminergic neurodegeneration in an AAV-synuclein mouse model of Parkinson's disease. Mol Neurodegener. 5: 42.
	5. Kovács I <i>et al.</i> (2015) Reactive oxygen species-mediated bacterial killing by B
	lymphocytes. <u>J Leukoc Biol. 97 (6): 1133-7.</u>
Storage	Store at +4°C or at -20°C if preferred.
	Storage in frost-free freezers is not recommended.
	This product should be stored undiluted. Avoid repeated freezing and thawing as this may
	denature the antibody. Should this product contain a precipitate we recommend
	microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety	Material Safety Datasheet documentation #10040 available at:
Information	https://www.bio-rad-antibodies.com/SDS/MCA4685
	10040
Regulatory	For research purposes only

## **Related Products**

# **Recommended Secondary Antibodies**

Rabbit Anti Mouse IgG (STAR12...)

Goat Anti Mouse IgG IgA IgM (STAR87...) HRP

Goat Anti Mouse IgG (STAR76...)

RPE

Goat Anti Mouse IgG (STAR70...)

FITC

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Rabbit Anti Mouse IgG (STAR9...) FITC

Goat Anti Mouse IgG (STAR77...) HRP

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

Rabbit Anti Mouse IgG (STAR13...) <u>HRP</u>

**Recommended Negative Controls** 

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

Fax: +1 919 878 3751 Fax: +44 (0)1865 852 739 Fax: +49 (0) 89 8090 95 50 To Email: antibody\_sales\_us@bio-rad.com Email: antibody\_sales\_uk@bio-rad.com Email: antibody\_sales\_de@bio-rad.com Email: antibody\_sales\_de@bio-rad.com

batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M367890:200529'

#### Printed on 20 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint