

Datasheet: MCA4685GA BATCH NUMBER 167503

Description:	MOUSE ANTI HUMAN CYTOCHROME B245 HEAVY CHAIN		
Specificity:	CYTOCHROME B245 HEAVY CHAIN		
Other names:	gp91phox		
Format:	Purified		
Product Type:	Monoclonal Antibody		
Clone:	NL7		
Isotype:	lgG1		
Quantity:	0.1 mg		

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .				
		Yes	No	Not Determined	Suggested Dilution
	Flow Cytometry (1)				1/25 - 1/200
	Immunohistology - Frozen				
	Immunohistology - Paraffin			•	
	ELISA	-			
	Immunoprecipitation				
	Western Blotting				
	Immunofluorescence	•			
	 Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls. (1) Membrane permeabilization is required for this application. The use of Leucoperm (Product Code <u>BUF09</u>) is recommended for this purpose. 				
Target Species	Human				
Product Form	Purified IgG - liquid				
Preparation	Purified IgG prepared by supernatant	affinity c	hromatog	raphy on Protein A fror	n tissue culture
Buffer Solution	Phosphate buffered salin	e			

Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)			
Carrier Free	Yes			
Approx. Protein Concentrations	IgG concentration 1.0mg/ml			
Immunogen	Cytochrome B245 solubilized in octyl-beta-glucopyranoside.			
External Database Links	UniProt: <u>P04839</u> <u>Related reagents</u> Entrez Gene: <u>1536</u> CYBB <u>Related reagents</u>			
Synonyms	NOX2			
RRID	AB_1963550			
Fusion Partners	Spleen cells from immunized BALB/c were fused with cells of the 3PU1 myeloma cell line.			
Specificity	 Mouse anti Human Cytochrome B245 Heavy Chain antibody, clone NL7 recognizes the heavy chain of cytochrome B245, also known as gp91phox or NOX2, a subunit of the nicotinamide adenine dinucleotide phosphate-oxidase (NADPH oxidase). NADPH oxidase is a membrane-bound enzyme complex in neutrophils that produces the superoxide anion (O₂⁻) by transferring single electrons from NADPH inside the cell across the membrane and coupling them to molecular oxygen. Superoxide is used for the destruction of pathogens. Individuals with defects in NADPH oxidase suffer from chronic granulomatous disease, characterised by severe recurrent infections. The heavy chain (gp91phox), along with the light chain (p22phox) form cytochrome B245 (also known as flavocytochome b, or Cyt b), which is always membrane-resident. Upon the appropriate stimulus, other components of NADPH oxidase (p47phox, p40phox, p67phox and Rac2) rapidly translocate from the cytosol to the cell membrane to form the complete enzyme complex. Cytochrome B245 appears to serve as a scaffold for the assembly of the other cytosolic units of the oxidase and to provide a transmembrane pathway for electrons. Mouse anti Human Cytochrome B245 Heavy Chain antibody, clone NL7 binds amino acids 498-506 (EKDVITGLK), a cytosolic epitope on gp91phox not accessible on intact neutrophils. It recognizes the native as well as the denatured form of the protein. Clone NL7 has an inhibitory effect on NADPH oxidase activation. Research indicates that this inhibitory effect is exerted during the initial stages of oxidase activation. The antibody does not interfere with the translocation of the cytosolic subunits of NADPH oxidase to the cell 			
Flow Cytometry	Les 10 μ of the suggested working dilution to lobal 1 \times 10 ⁶ calls in 100 μ			

Flow Cytometry

Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.

Western Blotting	MCA4685GA detects a broad band of approximately 90kDa
References	 Burritt, J.B. <i>et al.</i> (2003) Functional epitope on human neutrophil flavocytochrome b558. JImmunol. 170 (12): 6082-9. Taylor, R.M. <i>et al.</i> (2004) Site-specific inhibitors of NADPH oxidase activity and structural probes of flavocytochrome b: characterization of six monoclonal antibodies to the p22phox subunit. JImmunol. 173 (12): 7349-57. Taylor, R.M. <i>et al.</i> (2007) Cloning, sequence analysis and confirmation of derived gene sequences for three epitope-mapped monoclonal antibodies against human phagocyte flavocytochrome b. Mol Immunol. 44 (4): 625-37. Cao, S. <i>et al.</i> (2010) Fcγ receptors are required for NF-κB signaling, microglial activation and dopaminergic neurodegeneration in an AAV-synuclein mouse model of Parkinson's disease. Mol Neurodegener. 5: 42. Kovács I <i>et al.</i> (2015) Reactive oxygen species-mediated bacterial killing by B lymphocytes. J Leukoc Biol. 97 (6): 1133-7.
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA4685GA 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12)	RPE		
Goat Anti Mouse IgG IgA IgM (STAR87) <u>HRP</u>			
Goat Anti Mouse IgG (STAR76)	RPE		
Goat Anti Mouse IgG (STAR70)	<u>FITC</u>		
Goat Anti Mouse IgG (H/L) (STAR117)	<u>Alk. Phos., DyLight®488, DyLight®550,</u>		
	DyLight®650, DyLight®680, DyLight®800,		
	<u>FITC, HRP</u>		
Rabbit Anti Mouse IgG (STAR9)	FITC		
Goat Anti Mouse IgG (STAR77)	HRP		
Goat Anti Mouse IgG (Fc) (STAR120)	FITC, HRP		
Rabbit Anti Mouse IgG (STAR13)	HRP		
Recommended Negative Controls			

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	d.com	Email: antibody_sales_uk@bio-ra	ad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M429130:240320'

Printed on 20 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint