

## Datasheet: MCA469F

**BATCH NUMBER 165317**

<b>Description:</b>	MOUSE ANTI HUMAN CD9:FITC
<b>Specificity:</b>	CD9
<b>Other names:</b>	MRP-1
<b>Format:</b>	FITC
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	MM2/57
<b>Isotype:</b>	IgG2b
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

### Target Species

Human

### Species Cross Reactivity

Reacts with: Cat, Rhesus Monkey, Bovine, Dog, Rabbit, Horse, Pig, Mink, Llama, Ferret  
Based on sequence similarity, is expected to react with: Mustelid

**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

### Product Form

Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

### Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
FITC	490	525

### Preparation

Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant.

<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide 1% Bovine Serum Albumin
<b>Approx. Protein Concentrations</b>	IgG concentration 0.1 mg/ml
<b>Immunogen</b>	Human platelet membranes
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P21926</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">928</a>    CD9    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	MIC3, TSPAN29
<b>RRID</b>	AB_321492
<b>Fusion Partners</b>	Spleen cells from immunised BALB/c mice were fused with cells from the SP2/0 mouse myeloma line
<b>Specificity</b>	<p><b>Mouse anti Human CD9 antibody, clone MM2/57</b> recognizes human leukocyte antigen MIC3 also known as MRP-1 or CD9. CD9 is a 228 amino acid multi pass membrane glycoprotein belonging to the tetraspanin family with a molecular weight of ~24 kDa expressed by platelets, monocytes, some lymphocytes and endothelial cells.</p> <p>Mouse anti Human CD9 antibody, clone MM2/57 recognizes a conserved epitope on CD9 present on a wide range of mammalian species.</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
<b>References</b>	<ol style="list-style-type: none"> <li>1. Ed Knapp W. <i>et al.</i> (1989) Leucocyte Typing IV Oxford University Press.</li> <li>2. Jennings, L. K. <i>et al.</i> (1995) CD9 cluster workshop report: cell surface binding and functional analysis. In S.F. Sclossman. <i>et al.</i> Editors. 1995. Leucocyte Typing V. White Cell Differentiation Antigens. Oxford University Press, New York, NY. 1249-1251.</li> <li>3. Löffler, S. <i>et al.</i> (1997) CD9, a tetraspan transmembrane protein, renders cells susceptible to canine distemper virus. <a href="#">J Virol. 71: 42-9.</a></li> <li>4. Brodersen, R. <i>et al.</i> (1998) Analysis of the immunological cross reactivities of 213 well characterized monoclonal antibodies with specificities against various leucocyte surface antigens of human and 11 animal species. <a href="#">Vet Immunol Immunopathol. 64 (1): 1-13.</a></li> <li>5. Ferrer, M. <i>et al.</i> (1998) Pattern of expression of tetraspanin antigen genes in Burkitt lymphoma cell lines. <a href="#">Clin Exp Immunol. 113: 346-52.</a></li> <li>6. Kao, Y.R. <i>et al.</i> (2003) Tumor-associated antigen L6 and the invasion of human lung cancer cells. <a href="#">Clin Cancer Res. 9: 2807-16.</a></li> <li>7. Aasted, B. <i>et al.</i> (2007) Reactivity of monoclonal antibodies to human CD antigens with</li> </ol>

- cells from mink. [Vet Immunol Immunopathol. 119: 27-37.](#)
8. Davis, W.C. *et al.* (2007) Use of flow cytometry to identify monoclonal antibodies that recognize conserved epitopes on orthologous leukocyte differentiation antigens in goats, llamas, and rabbits. [Vet Immunol Immunopathol. 119: 123-30.](#)
  9. Meister, R.K. *et al.* (2007) Progress in the discovery and definition of monoclonal antibodies for use in feline research. [Vet Immunol Immunopathol. 119: 38-46.](#)
  10. Martel, C.J. & Aasted, B. (2009) Characterization of antibodies against ferret immunoglobulins, cytokines and CD markers. [Vet Immunol Immunopathol. 132:109-15.](#)
  11. Müller, T. *et al.* (2009) A novel embryonic stem cell line derived from the common marmoset monkey (*Callithrix jacchus*) exhibiting germ cell-like characteristics. [Hum Reprod. 24: 1359-72.](#)
  12. Kubota, H. *et al.* (2011) Glial cell line-derived neurotrophic factor and endothelial cells promote self-renewal of rabbit germ cells with spermatogonial stem cell properties. [FASEB J. 25 \(8\): 2604-14.](#)
  13. Hogue, I.B. *et al.* (2011) Gag induces the coalescence of clustered lipid rafts and tetraspanin-enriched microdomains at HIV-1 assembly sites on the plasma membrane. [J Virol. 85 \(19\): 9749-66.](#)
  14. Viswanathan, K. *et al.* (2017) Quantitative membrane proteomics reveals a role for tetraspanin enriched microdomains during entry of human cytomegalovirus. [PLoS One. 12 \(11\): e0187899.](#)
  15. Bearden, R.N. *et al.* (2017) *In-vitro* characterization of canine multipotent stromal cells isolated from synovium, bone marrow, and adipose tissue: a donor-matched comparative study. [Stem Cell Res Ther. 8 \(1\): 218.](#)
  16. Jackson, C.E. *et al.* (2017) Effects of Inhibiting VPS4 Support a General Role for ESCRTs in Extracellular Vesicle Biogenesis. [Biophys J. 113 \(6\): 1342-1352.](#)
  17. Fish, E.J. *et al.* (2018) Malignant canine mammary epithelial cells shed exosomes containing differentially expressed microRNA that regulate oncogenic networks. [BMC Cancer. 18 \(1\): 832.](#)
  18. Wąchalska, M. *et al.* (2020) Palmitoylated mNeonGreen Protein as a Tool for Visualization and Uptake Studies of Extracellular Vesicles. [Membranes \(Basel\). 10 \(12\): 373.](#)
  19. Fu, T. *et al.* (2021) Biomimetic vascularized adipose-derived mesenchymal stem cells bone-periosteum graft enhances angiogenesis and osteogenesis in a rabbit spine fusion model. [Stem Cell Res Therapy. 15 Sep Preprint \[Epub ahead of print\].](#)
  20. Kwasnik, M. *et al.* (2023) Protein-Coding Region Derived Small RNA in Exosomes from Influenza A Virus-Infected Cells. [Int J Mol Sci. 24 \(1\): 867.](#)

---

**Storage**

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

---

**Guarantee**

12 months from date of despatch

---

**Health And Safety**

Material Safety Datasheet documentation #10041 available at:

**Information** <https://www.bio-rad-antibodies.com/SDS/MCA469F10041>

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Negative Controls

[MOUSE IgG2b NEGATIVE CONTROL:FITC \(MCA691F\)](#)

### Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)  
'M385731:210513'

**Printed on 20 Oct 2023**

---

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)