

Datasheet: MCA46G

Description:	MOUSE ANTI RAT MHC CLASS II RT1B
Specificity:	MHC CLASS II RT1B
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	OX-6
Isotype:	IgG1
Quantity:	1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin (1)	▪			1/50 - 1/100
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose. PLP fixation is recommended for optimal results.

Target Species	Rat
Species Cross Reactivity	<p>Reacts with: Mouse</p> <p>N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
Product Form	Purified IgG - liquid

Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Rat thymocyte membrane glycoproteins.
RRID	AB_322113
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells from the NS1 mouse myeloma cell line.
Specificity	<p>Mouse anti Rat MHC Class II RT1B antibody, clone OX-6 recognizes a monomorphic determinant of the rat RT1B MHC class II antigen present on B lymphocytes, dendritic cells, some macrophages and certain epithelial cells.</p> <p>Rat MHC Class II RT1B antibody, clone OX-6 does not react with the rat BDIX strain due to a defect in RT1B expression (Male et al. 1987).</p> <p>The major histocompatibility complex (MHC) is a cluster of genes that are important in the immune response to infections. In rats, this complex is referred to as the RT1 region. In mice, this complex is referred to as the H-2 region.</p> <p>Mouse anti Rat MHC Class II RT1B antibody, clone OX-6 also cross reacts with a polymorphic determinant on mouse strains of the H-2 haplotypes k and s. Analysis of recombinant mouse strains has mapped the OX-6 determinant to the H-2I-A region (McMaster and Williams 1979 and Male et al. 1987).</p> <p>Mouse anti Rat MHC Class II RT1B antibody, clone OX-6 is routinely tested in flow cytometry on rat splenocytes.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> 1. Fernandez, J.L. & Weeks, M. (1986) Genetic monitoring of inbred strains of mice using monoclonal antibodies to major histocompatibility haplotypes and lymphocyte alloantigens. Lab Anim. 20 (4): 293-7. 2. Charteris, D.G. & Lightman, S.L. (1993) In vivo lymphokine production in experimental autoimmune uveoretinitis. Immunology. 78 (3): 387-92. 3. Whiteland, J.L. et al. (1995) Immunohistochemical detection of T-cell subsets and other leukocytes in paraffin-embedded rat and mouse tissues with monoclonal antibodies. J

[Histochem Cytochem. 43 \(3\): 313-20.](#)

4. McKechnie, N.M. *et al.* (1997) Immunization with the cross-reactive antigens Ov39 from *Onchocerca volvulus* and hr44 from human retinal tissue induces ocular pathology and activates retinal microglia. [J Infect Dis. 176 \(5\): 1334-43.](#)

5. Male, D.K. *et al.* (1987) Serological evidence for a defect in RT1.B (I-A) expression by the BDIX rat strain. [J Immunogenet. 14 \(6\): 301-12.](#)

6. Burrows, G.G. *et al.* (1998) Two-domain MHC class II molecules form stable complexes with myelin basic protein 69-89 peptide that detect and inhibit rat encephalitogenic T cells and treat experimental autoimmune encephalomyelitis. [J Immunol. 161 \(11\): 5987-96.](#)

7. Hofmann, N. *et al.* (2002) Increased expression of ICAM-1, VCAM-1, MCP-1, and MIP-1 alpha by spinal perivascular macrophages during experimental allergic encephalomyelitis in rats. [BMC Immunol. 3:11.](#)

8. Banerjee, S. *et al.* (2003) Development of organised conjunctival leucocyte aggregates after corneal transplantation in rats. [Br J Ophthalmol. 87: 1515-22.](#)

9. King, G.D. *et al.* (2008) Flt3L in combination with HSV1-TK-mediated gene therapy reverses brain tumor-induced behavioral deficits. [Mol Ther. 16: 682-90.](#)

10. Bode, U. *et al.* (2008) Dendritic cell subsets in lymph nodes are characterized by the specific draining area and influence the phenotype and fate of primed T cells. [Immunology. 123 \(4\): 480-90.](#)

11. Baca Jones, C.C. *et al.* (2009) Rat cytomegalovirus infection depletes MHC II in bone marrow derived dendritic cells. [Virology. 388: 78-90.](#)

12. Wang, Q. *et al.* (2009) Pyruvate protects against experimental stroke via an anti-inflammatory mechanism. [Neurobiol Dis. 2009 Oct;36\(1\):223-31.](#)

13. Zilka, N. *et al.* (2009) Human misfolded truncated tau protein promotes activation of microglia and leukocyte infiltration in the transgenic rat model of tauopathy. [J Neuroimmunol. 209 \(1-2\): 16-25.](#)

14. Calvo, M. *et al.* (2010) Neuregulin-ErbB signaling promotes microglial proliferation and chemotaxis contributing to microgliosis and pain after peripheral nerve injury. [J Neurosci. 30 \(15\): 5437-50.](#)

15. Bereczky-Veress, B. *et al.* (2010) Influence of perineurial cells and Toll-like receptors 2 and 9 on Herpes simplex type 1 entry to the central nervous system in rat encephalitis. [PLoS One. 5\(8\): e12350.](#)

16. Kawamura, J. *et al.* (2010) Neuron-immune Interactions in the Sensitized Thalamus Induced by Mustard Oil Application to Rat Molar Pulp. [J Dent Res. 89: 1309-14.](#)

17. McClain, J.A. *et al.* (2011) Adolescent binge alcohol exposure induces long-lasting partial activation of microglia. [Brain Behav Immun. 25 Suppl 1: S120-8.](#)

18. Lobato-Pascual, A. *et al.* (2013) Rat macrophage C-type lectin is an activating receptor expressed by phagocytic cells. [PLoS One. 8: e57406.](#)

19. Maneu, V. *et al.* (2016) Immunosuppression, peripheral inflammation and invasive infection from endogenous gut microbiota activate retinal microglia in mouse models. [Microbiol Immunol. 60 \(9\): 617-25.](#)

20. Collins, J.J.P. *et al.* (2018) Impaired Angiogenic Supportive Capacity and Altered Gene Expression Profile of Resident CD146⁺ Mesenchymal Stromal Cells Isolated from Hyperoxia-Injured Neonatal Rat Lungs. [Stem Cells Dev. 27 \(16\): 1109-24.](#)

21. Takizawa, T. *et al.* (2017) High-mobility group box 1 is an important mediator of microglial activation induced by cortical spreading depression. [J Cereb Blood Flow Metab. 37 \(3\): 890-901.](#)

22. Liu, M. *et al.* (2017) Pioglitazone Attenuates Neuroinflammation and Promotes Dopaminergic Neuronal Survival in the Nigrostriatal System of Rats after Diffuse Brain Injury. [J Neurotrauma. 34 \(2\): 414-22.](#)
23. Noailles, A. *et al.* (2018) Systemic inflammation induced by lipopolysaccharide aggravates inherited retinal dystrophy. [Cell Death Dis. 9 \(3\): 350.](#)
24. Stangl, H. *et al.* (2020) MHC/class-II-positive cells inhibit corticosterone of adrenal gland cells in experimental arthritis: a role for IL-1 β , IL-18, and the inflammasome. [Sci Rep. 10 \(1\): 17071.](#)
25. Silva, B.A. *et al.* (2022) Understanding the role of the blood brain barrier and peripheral inflammation on behavior and pathology on ongoing confined cortical lesions. [Mult Scler Relat Disord. 57: 103346.](#)
26. Sawanobori, Y. *et al.* (2021) Selective involution of thymic medulla by cyclosporine A with a decrease of mature thymic epithelia, XCR1⁺ dendritic cells, and epithelium-free areas containing Foxp3⁺ thymic regulatory T cells. [Histochem Cell Biol. 156 \(2\): 133-146.](#)
27. Koppe, C. *et al.* (2021) Local Inflammatory Response after Intramuscularly Implantation of Anti-Adhesive Plasma-Fluorocarbon-Polymer Coated Ti6Al4V Discs in Rats. [Polymers \(Basel\). 13 \(16\): 2684.](#)
28. Matsuyama, S. *et al.* (2021) Properties of macrophages and lymphocytes appearing in rat renal fibrosis followed by repeated injection of cisplatin. [J Vet Med Sci. 83 \(9\): 1435-42.](#)
29. Sarkar, T. *et al.* (2022) Perinatal exposure to synergistic multiple stressors leads to cellular and behavioral deficits mimicking Schizophrenia-like pathology. [Biol Open. 11\(3\):bio058870.](#)
30. Pervin, M. *et al.* (2022) Possible Cytoprotection of Low Dose Lipopolysaccharide in Rat Thioacetamide-Induced Liver Lesions, Focusing on the Analyses of Hepatic Macrophages and Autophagy. [Toxicol Pathol. : 1926233221076758.](#)
31. akala-Jakimowicz, M. & Puzianowska-Kuznicka, M. (2022) Towards Understanding the Lymph Node Response to Skin Infection with Saprophytic Staphylococcus epidermidis. [Biomedicines. 10\(5\):1021.](#)
32. Zakani, M. *et al.* (2023) Paths to hippocampal damage in neuromyelitis optica spectrum disorders [Neuropathol App Neurobiol. 49 \(2\) e12893](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA46G>
10040

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA1209\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------	------------------	-------------------------------------------------------------------------------------------------------------------------------------------------	---------------	-----------------------------------------------------------------------------------------------------------------------------------------------------

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M384088:210513'

Printed on 15 Aug 2023