

## Datasheet: MCA46R

**BATCH NUMBER 1114**

<b>Description:</b>	MOUSE ANTI RAT MHC CLASS II RT1B
<b>Specificity:</b>	MHC CLASS II RT1B
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	OX-6
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.25 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin (1)	▪			1/50 - 1/100
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

**(1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose. PLP fixation is recommended for optimal results.**

<b>Target Species</b>	Rat
<b>Species Cross Reactivity</b>	<p>Reacts with: Mouse</p> <p><b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>

<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide
<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Rat thymocyte membrane glycoproteins.
<b>RRID</b>	AB_322404
<b>Fusion Partners</b>	Spleen cells from immunised BALB/c mice were fused with cells from the NS1 mouse myeloma cell line.
<b>Specificity</b>	<p><b>Mouse anti Rat MHC Class II RT1B antibody, clone OX-6</b> recognizes a monomorphic determinant of the rat RT1B MHC class II antigen present on B lymphocytes, dendritic cells, some macrophages and certain epithelial cells.</p> <p>Rat MHC Class II RT1B antibody, clone OX-6 does not react with the rat BDIX strain due to a defect in RT1B expression (<a href="#">Male et al. 1987</a>).</p> <p>The major histocompatibility complex (MHC) is a cluster of genes that are important in the immune response to infections. In rats, this complex is referred to as the RT1 region. In mice, this complex is referred to as the H-2 region.</p> <p>Mouse anti Rat MHC Class II RT1B antibody, clone OX-6 also cross reacts with a polymorphic determinant on mouse strains of the H-2 haplotypes k and s. Analysis of recombinant mouse strains has mapped the OX-6 determinant to the H-2I-A region (<a href="#">McMaster and Williams 1979</a> and <a href="#">Male et al. 1987</a>).</p> <p>Mouse anti Rat MHC Class II RT1B antibody, clone OX-6 is routinely tested in flow cytometry on rat splenocytes.</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label $10^6$ cells in 100ul.
<b>References</b>	<ol style="list-style-type: none"> <li>McMaster, W.R. &amp; Williams, A.F. (1979) Identification of Ia glycoproteins in rat thymus and purification from rat spleen. <a href="#">Eur J Immunol. 9 (6): 426-33.</a></li> <li>Fernandez, J.L. &amp; Weeks, M. (1986) Genetic monitoring of inbred strains of mice using monoclonal antibodies to major histocompatibility haplotypes and lymphocyte alloantigens. <a href="#">Lab Anim. 20 (4): 293-7.</a></li> <li>Charteris, D.G. &amp; Lightman, S.L. (1993) In vivo lymphokine production in experimental</li> </ol>

- autoimmune uveoretinitis. [Immunology. 78 \(3\): 387-92.](#)
4. Whiteland, J.L. *et al.* (1995) Immunohistochemical detection of T-cell subsets and other leukocytes in paraffin-embedded rat and mouse tissues with monoclonal antibodies. [J Histochem Cytochem. 43 \(3\): 313-20.](#)
  5. McKechnie, N.M. *et al.* (1997) Immunization with the cross-reactive antigens Ov39 from *Onchocerca volvulus* and hr44 from human retinal tissue induces ocular pathology and activates retinal microglia. [J Infect Dis. 176 \(5\): 1334-43.](#)
  6. Male, D.K. *et al.* (1987) Serological evidence for a defect in RT1.B (I-A) expression by the BDIX rat strain. [J Immunogenet. 14 \(6\): 301-12.](#)
  7. Burrows, G.G. *et al.* (1998) Two-domain MHC class II molecules form stable complexes with myelin basic protein 69-89 peptide that detect and inhibit rat encephalitogenic T cells and treat experimental autoimmune encephalomyelitis. [J Immunol. 161 \(11\): 5987-96.](#)
  8. Zilka, N. *et al.* (2009) Human misfolded truncated tau protein promotes activation of microglia and leukocyte infiltration in the transgenic rat model of tauopathy. [J Neuroimmunol. 209 \(1-2\): 16-25.](#)
  9. Kawamura, J. *et al.* (2010) Neuron-immune Interactions in the Sensitized Thalamus Induced by Mustard Oil Application to Rat Molar Pulp. [J Dent Res. 89: 1309-14.](#)
  10. Calvo, M. *et al.* (2010) Neuregulin-ErbB signaling promotes microglial proliferation and chemotaxis contributing to microgliosis and pain after peripheral nerve injury. [J Neurosci. 30 \(15\): 5437-50.](#)
  11. McClain, J.A. *et al.* (2011) Adolescent binge alcohol exposure induces long-lasting partial activation of microglia. [Brain Behav Immun. 25 Suppl 1: S120-8.](#)
  12. Baca Jones, C.C. *et al.* (2009) Rat cytomegalovirus infection depletes MHC II in bone marrow derived dendritic cells. [Virology. 388: 78-90.](#)
  13. Lobato-Pascual, A. *et al.* (2013) Rat macrophage C-type lectin is an activating receptor expressed by phagocytic cells. [PLoS One. 8: e57406.](#)
  14. Takizawa, T. *et al.* (2016) High-mobility group box 1 is an important mediator of microglial activation induced by cortical spreading depression [Journal of Cerebral Blood Flow & Metabolism. May 3 \[Epub ahead of print\]](#)
  15. Liu, M. *et al.* (2017) Pioglitazone Attenuates Neuroinflammation and Promotes Dopaminergic Neuronal Survival in the Nigrostriatal System of Rats after Diffuse Brain Injury. [J Neurotrauma. 34 \(2\): 414-22.](#)
  16. Noailles, A. *et al.* (2018) Systemic inflammation induced by lipopolysaccharide aggravates inherited retinal dystrophy. [Cell Death Dis. 9 \(3\): 350.](#)
  17. Stangl, H. *et al.* (2020) MHC/class-II-positive cells inhibit corticosterone of adrenal gland cells in experimental arthritis: a role for IL-1 $\beta$ , IL-18, and the inflammasome. [Sci Rep. 10 \(1\): 17071.](#)

---

#### Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

#### Guarantee

12 months from date of despatch

---

**Health And Safety Information**      Material Safety Datasheet documentation #10040 available at:  
<https://www.bio-rad-antibodies.com/SDS/MCA46R10040>

**Regulatory**      For research purposes only

## Related Products

### Recommended Secondary Antibodies

- Rabbit Anti Mouse IgG (STAR12...)      [RPE](#)  
Goat Anti Mouse IgG IgA IgM (STAR87...) [HRP](#)  
Goat Anti Mouse IgG (STAR76...)      [RPE](#)  
Goat Anti Mouse IgG (STAR70...)      [FITC](#)  
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),  
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),  
[FITC](#), [HRP](#)  
  
Rabbit Anti Mouse IgG (STAR9...)      [FITC](#)  
Rabbit Anti Mouse IgG (STAR13...)      [HRP](#)  
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)  
Goat Anti Mouse IgG (STAR77...)      [HRP](#)

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA1209\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)  
'M367936:200529'

Printed on 15 Aug 2024