Datasheet: MCA4713 BATCH NUMBER 159478

Description:	RAT ANTI MOUSE P2X7
Specificity:	P2X7
Other names:	P2RX7
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	Hano43
Isotype:	lgG2b
Quantity:	0.25 mg

Product Details

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u>.

Yes	No	Not Determined	Suggested Dilution
-			1/25 - 1/200
		-	
		•	
		•	
		-	
		•	
	•		•

necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative	0.09% Sodium Azide (NaN ₃)

Stabilisers

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
Immunogen	A P2X7-expression construct and a final boost with P2X7-transfected HEK cells.
External Database Links	UniProt: <u>Q9Z1M0</u> Related reagents
	Entrez Gene:
	18439 P2rx7 Related reagents
Synonyms	P2x7
RRID	AB_10545124
Fusion Partners	Spleen cells from immunised rats were fused with cells of the Sp2/0 myeloma cell line.
Specificity	Rat anti Mouse P2X7 antibody, clone Hano43 recognizes the P2X purinoceptor 7, also known as P2X7. The P2X7 ATP receptor has a distinctive long C-terminal tail with multiple potential protein and lipid interaction motifs and is highly polymorphic. It is a cation selective ion channel that opens up on binding of extracellular ATP. Sustained activation by extracellular ATP results in the formation of a reversible pore in the plasma membrane that is permeable to hydrophilic solutes of up to 900 Da. Once a pore is opened massive upset of cytoplasmic ion homeostasis occurs and the pore stays open as long as it is bound by ATP. Should ATP stimulation continue the cell will become irreversibly damaged and die. P2X7 plays a key role in the maturation and release of IL-1 and other IL-1 family members during inflammation. As such, P2X7 blockers might be useful as anti-inflammatory agents.
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.
References	 Adriouch, S. <i>et al.</i> (2005) Probing the expression and function of the P2X7 purinoceptor with antibodies raised by genetic immunization. <u>Cell Immunol. 236: 72-7.</u> Adriouch, S. <i>et al.</i> (2008) ADP-ribosylation at R125 gates the P2X7 ion channel by presenting a covalent ligand to its nucleotide binding site. <u>FASEB J. 2008</u> <u>Mar:22(3):861-9.</u> Aswad, F. and Dennert, G. (2006) P2X7 receptor expression levels determine lethal effects of a purine based danger signal in T lymphocytes. <u>Cell Immunol. 243: 58-65.</u> Adriouch, S. <i>et al.</i> (2009) Characterisation of the R276A gain-of-function mutation in the ectodomain of murine P2X7. <u>Purinergic Signal. 5: 151-61.</u> Kurashima, Y. <i>et al.</i> (2012) Extracellular ATP mediates mast cell-dependent intestinal inflammation through P2X7 purinoceptors. <u>Nat Commun. 3: 1034.</u> Hu, S.J. <i>et al.</i> (2015) Upregulation of P2RX7 in Cx3cr1-Deficient Mononuclear Phagocytes Leads to Increased Interleukin-1β Secretion and Photoreceptor

	 Neurodegeneration. <u>J Neurosci. 35 (18): 6987-96.</u> 7. Barabási, B. <i>et al.</i> (2016) Effect of axotomy and 17β-estradiol on P2X7 receptor expression pattern in the hypoglossal nucleus of ovariectomized mice. <u>Neuroscience. 319: 107-15.</u> 8. Liu, Q. & Kim, C.H. (2019) Control of Tissue-Resident Invariant NKT Cells by Vitamin A Metabolites and P2X7-Mediated Cell Death. <u>J Immunol. 203 (5): 1189-97.</u>
Further Reading	 Ferrari, D. <i>et al.</i> (2006) The P2X7 receptor: a key player in IL-1 processing and release. <u>J Immunol. 176: 3877-83.</u> Schwarz, N. <i>et al</i> (2009) Activation of the P2X7 ion channel by soluble and covalently bound ligands. <u>Purinergic Signal. 5: 139-49.</u>
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA4713 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Rat IgG (STAR69)		<u>FITC</u>				
Goat Anti Rat IgG (STAR73)		RPE				
Goat Ant	i Rat IgG (MOUSE ADSC	ORBED) (STA	AR71) <u>DyLight®550</u> ,	DyLight®650,	DyLight®800	
Rabbit Ar	nti Rat IgG (STAR21)		HRP			
Rabbit Anti Rat IgG (STAR16)			DyLight®800			
Goat Anti Rat IgG (STAR131)			<u>Alk. Phos., Bic</u>	otin		
Rabbit Ar	nti Rat IgG (STAR17)		<u>FITC</u>			
Goat Ant	i Rat IgG (STAR72)		HRP			
North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21	
America	Fax: +1 919 878 3751	Wondwide	Fax: +44 (0)1865 852 739	Latope	Fax: +49 (0) 89 8090 95 50	

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M384230:210513'

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint

Email: antibody_sales_us@bio-rad.com