

Datasheet: MCA4713A488

Description:	RAT ANTI MOUSE P2X7:Alexa Fluor® 488		
Specificity:	P2X7		
Other names:	P2RX7		
Format:	ALEXA FLUOR® 488		
Product Type:	Monoclonal Antibody		
Clone:	Hano43		
Isotype:	lgG2b		
Quantity:	100 TESTS/1ml		

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				Neat
Immunohistology - Frozen			•	
Immunohistology - Paraffin			•	
ELISA			•	
Immunoprecipitation			•	
Western Blotting			•	
Immunofluorescence				

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse				
Product Form	Purified IgG conjugated to Alexa Fluor 488 - liquid				
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)		
	Alexa Fluor®488	495	519		
Preparation	Purified IgG prepar supernatant	red by affinity chromatog	raphy on Protein G t		
Buffer Solution	Phosphate buffered	d saline			

Preservative	0.09% Sodium Azide (NaN ₃)				
Stabilisers	1% Bovine Serum Albumin				
Approx. Protein Concentrations	IgG concentration 0.05mg/ml				
Immunogen	A P2X7-expression construct and a final boost with P2X7-transfected HEK cells.				
External Database					
Links	UniProt:				
	Q9Z1M0 Related reagents				
	Entrez Gene:				
	18439 P2rx7 Related reagents				
Synonyms	P2x7				
RRID	AB_2158252				
Fusion Partners	Spleen cells from immunised rats were fused with cells of the Sp2/0 myeloma cell line.				
Specificity	Rat anti Mouse P2X7 antibody, clone Hano43 recognizes the P2X purinoceptor 7, also known as P2X7. The P2X7 ATP receptor has a distinctive long C-terminal tail with multiple potential protein and lipid interaction motifs and is highly polymorphic. It is a cation selective ion channel that opens up on binding of extracellular ATP. Sustained activation by extracellular ATP results in the formation of a reversible pore in the plasma membrane that is permeable to hydrophilic solutes of up to 900 Da. Once a pore is opened massive upset of cytoplasmic ion homeostasis occurs and the pore stays open as long as it is bound by ATP. Should ATP stimulation continue the cell will become irreversibly damaged and die. P2X7 plays a key role in the maturation and release of IL-1 and other IL-1 family members during inflammation. As such, P2X7 blockers might be useful as anti-inflammatory agents.				
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.				
References	 Adriouch, S. <i>et al.</i> (2005) Probing the expression and function of the P2X7 purinoceptor with antibodies raised by genetic immunization. <u>Cell Immunol. 236: 72-7.</u> Adriouch, S. <i>et al.</i> (2008) ADP-ribosylation at R125 gates the P2X7 ion channel by presenting a covalent ligand to its nucleotide binding site. <u>FASEB J. 2008</u> <u>Mar;22(3):861-9.</u> Aswad, F. and Dennert, G. (2006) P2X7 receptor expression levels determine lethal effects of a purine based danger signal in T lymphocytes. <u>Cell Immunol. 243: 58-65.</u> Adriouch, S. <i>et al.</i> (2009) Characterisation of the R276A gain-of-function mutation in the ectodomain of murine P2X7. <u>Purinergic Signal. 5: 151-61.</u> Kurashima, Y. <i>et al.</i> (2012) Extracellular ATP mediates mast cell-dependent intestinal 				

- 5. Kurashima, Y. et al. (2012) Extracellular ATP mediates mast cell-dependent intestinal inflammation through P2X7 purinoceptors. Nat Commun. 3: 1034.
- 6. Hu, S.J. et al. (2015) Upregulation of P2RX7 in Cx3cr1-Deficient Mononuclear Phagocytes Leads to Increased Interleukin-1 β Secretion and Photoreceptor

Neurodegeneration. J Neurosci. 35 (18): 6987-96.

- 7. Barabási, B. et al. (2016) Effect of axotomy and 17β-estradiol on P2X7 receptor expression pattern in the hypoglossal nucleus of ovariectomized mice. Neuroscience. 319: 107-15.
- 8. Liu, Q. & Kim, C.H. (2019) Control of Tissue-Resident Invariant NKT Cells by Vitamin A Metabolites and P2X7-Mediated Cell Death. J Immunol. 203 (5): 1189-97.

Further Reading

- 1. Ferrari, D. et al. (2006) The P2X7 receptor: a key player in IL-1 processing and release. J Immunol. 176: 3877-83.
- 2. Schwarz, N. et al (2009) Activation of the P2X7 ion channel by soluble and covalently bound ligands. Purinergic Signal. 5: 139-49.

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee

12 months from date of despatch

Acknowledgements

This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

Health And Safety Information

Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA4713A488 10041

Regulatory

For research purposes only

Related Products

Recommended Useful Reagents

MOUSE SEROBLOCK FcR (BUF041A) MOUSE SEROBLOCK FcR (BUF041B)

America

North & South Tel: +1 800 265 7376 Fax: +1 919 878 3751 Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

Printed on 12 Aug 2023

© 2023 Bio-Rad Laboratories Inc | Legal | Imprint