

Datasheet: MCA4738

Description:	MOUSE ANTI CLOSTRIDIUM DIFFICILE TOXIN B
Specificity:	CLOSTRIDIUM DIFFICILE TOXIN B
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	5158
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

RRID AB_1658138

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			1/20 - 1/200
Immunoprecipitation			▪	
Western Blotting			▪	
Immunofluorescence			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species Bacterial

Product Form Purified IgG - liquid

Preparation Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

Buffer Solution Phosphate buffered saline

Preservative Stabilisers 0.1% Sodium Azide (NaN₃)

Approx. Protein Concentrations IgG concentration 0.1mg/ml

External Database Links

UniProt:

[P18177](#)

[Related reagents](#)

Synonyms	tcdB
Specificity	<p>Mouse anti Clostridium difficile Toxin B antibody, clone 5158 recognizes <i>Clostridium difficile</i> toxin B. No reaction is observed with toxin A. <i>Clostridium difficile</i> is a gram-positive motile bacterium which is the leading cause of diarrhoea in developed countries. The incidence of disease in humans varies greatly with age, spore density and the administration of antibiotics.</p> <p>Toxin B (TcdB) is a large exotoxin. Its role in disease is less well understood than that of toxin A, though it seems clear that TcdB contributes to disease. Both toxins modify the Ras superfamily of small GTPases via glycosylation, inactivating them and leading to the disruption of vital signaling pathways in the cell.</p>
Further Reading	1. Voth, D.E. & Ballard, J.D. (2005) Clostridium difficile toxins: mechanism of action and role in disease. Clin Microbiol Rev. 18 (2): 247-63.
Storage	<p>Store at +4°C or at -20°C if preferred.</p> <p>Storage in frost-free freezers is not recommended.</p> <p>This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
Guarantee	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Rabbit Anti Mouse IgG (STAR8...)	DyLight@800
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight@488 , DyLight@680 , DyLight@800 , FITC , HRP

North & South America Tel: +1 800 265 7376
 Fax: +1 919 878 3751
 Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
 Fax: +44 (0)1865 852 739
 Email: antibody_sales_uk@bio-rad.com

Europe Tel: +49 (0) 89 8090 95 21
 Fax: +49 (0) 89 8090 95 50
 Email: antibody_sales_de@bio-rad.com

'M358002:190911'

Printed on 11 Oct 2019

