

Datasheet: MCA4754A647

BATCH NUMBER 173949

| | |
|----------------------|-------------------------------------|
| Description: | MOUSE ANTI RAT CD63:Alexa Fluor®647 |
| Specificity: | CD63 |
| Format: | ALEXA FLUOR® 647 |
| Product Type: | Monoclonal Antibody |
| Clone: | AD1 |
| Isotype: | IgG1 |
| Quantity: | 100 TESTS/1ml |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------|-----|----|----------------|--------------------|
| Flow Cytometry | ▪ | | | Neat - 1/10 |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

| | | | |
|---------------------------------------|---|----------------------------|--------------------------|
| Target Species | Rat | | |
| Product Form | Purified IgG conjugated to Alexa Fluor® 647 - liquid | | |
| Max Ex/Em | Fluorophore | Excitation Max (nm) | Emission Max (nm) |
| | Alexa Fluor®647 | 650 | 665 |
| Preparation | Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant | | |
| Buffer Solution | Phosphate buffered saline | | |
| Preservative | 0.09% Sodium Azide (NaN ₃) | | |
| Stabilisers | 1% Bovine Serum Albumin | | |
| Approx. Protein Concentrations | IgG concentration 0.05mg/ml | | |

| | |
|--------------------------------|--|
| Immunogen | RBL-2H3 rat basophilic leukaemia cell line. |
| External Database Links | <p>UniProt: P28648 Related reagents</p> <p>Entrez Gene: 29186 Cd63 Related reagents</p> |
| RRID | AB_10613490 |
| Fusion Partners | Spleen cells from immunized BALB/c mice were fused with cells of the X63-Ag8-653 myeloma cell line. |
| Specificity | <p>Mouse anti Rat CD63 antibody, clone AD1 recognizes rat CD63, an intracellular type-III lysosomal glycoprotein and member of the tetraspanin (TM4SF) family, found on activated platelets and resident macrophages. Rat CD63 is expressed on mast cells in tissues and can be induced on other cell types <i>in vitro</i> (Nishikata et al. 1992).</p> <p>Mouse anti Rat CD63 antibody, clone AD1 partially inhibits IgE-mediated histamine release and binding competition assays indicate that it is sterically situated close to the high-affinity IgE receptor on rat basophilic leukemia (RBL-2H3) cells (Kitani et al. 1991).</p> |
| Flow Cytometry | Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul. |
| References | <ol style="list-style-type: none"> 1. Boros, M. <i>et al.</i> (1995) Response of mucosal mast cells to intestinal ischemia-reperfusion injury in the rat. Shock. 3 (2): 125-31. 2. Vincent-Schneider, H. <i>et al.</i> (2001) Secretory granules of mast cells accumulate mature and immature MHC class II molecules J Cell Sci. 114(Pt 2): 323-34. 3. Kraft, S. <i>et al.</i> (2005) Anti-CD63 antibodies suppress IgE-dependent allergic reactions <i>in vitro</i> and <i>in vivo</i>. J Exp Med. 201: 385-96. 4. Ramezani-Moghadam, M. <i>et al.</i> (2015) Adiponectin Reduces Hepatic Stellate Cell Migration by Promoting Tissue Inhibitor of Metalloproteinase-1 (TIMP-1) Secretion. J Biol Chem. 290(9): 5533-42. 5. Yoshimura, A. <i>et al.</i> (2016) Generation of a novel transgenic rat model for tracing extracellular vesicles in body fluids. Sci Rep. 6: 31172. 6. Pasquier, A. <i>et al.</i> (2019) Lysosomal degradation of newly formed insulin granules contributes to β cell failure in diabetes. Nat Commun. 10 (1): 3312. 7. Pusic, K.M. <i>et al.</i> (2019) IFNγ-Stimulated Dendritic Cell Exosomes for Treatment of Migraine Modeled Using Spreading Depression. Front Neurosci. 13: 942. |
| Storage | <p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.</p> |

Guarantee 12 months from date of despatch

Acknowledgements This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

Health And Safety Information Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA4754A647>

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:Alexa Fluor® 647 \(MCA1209A647\)](#)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M385514:210513'

Printed on 27 May 2026