

## Datasheet: MCA4770GA

<b>Description:</b>	MOUSE ANTI HUMAN TALIN-1
<b>Specificity:</b>	TALIN-1
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	97H6
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin (1)	▪			1/25 - 1/200
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting	▪			
Immunofluorescence (2)	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

**(1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.**

**(2) Methanol treatment is required prior to antibody staining**

<b>Target Species</b>	Human
<b>Species Cross Reactivity</b>	Reacts with: Mouse, Rabbit <b>N.B.</b> Antibody reactivity and working conditions may vary between species.
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> )
<b>Carrier Free</b>	Yes

<b>Approx. Protein Concentrations</b>	IgG concentration 1.0mg/ml
<b>Immunogen</b>	his-tagged polypeptide containing residues 489–911 of mouse talin1.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">Q9Y490</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">7094</a>    TLN1    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	KIAA1027, TLN
<b>RRID</b>	AB_10675618
<b>Fusion Partners</b>	Spleen cells from immunized Balb/c mice were fused with cells from the NS0 murine myeloma.
<b>Specificity</b>	<b>Mouse anti Human Talin-1 antibody, clone 97H6</b> recognizes an epitope within the region of amino acids <a href="#">482-655</a> of human Talin-1, a high molecular weight cytoskeletal protein, linking vinculin to the integrins, thereby linking the cytoskeleton to the extracellular matrix. It is specifically found in regions of cell-substratum contact and in lymphocytes at cell-cell contact points. Talin-1 plays a role in the assembly of actin filaments and in the migration of cells such as osteoclasts and fibroblasts.
<b>Histology Positive Control Tissue</b>	Tonsil
<b>References</b>	<ol style="list-style-type: none"> <li>1. Kopp, P.M. <i>et al.</i> (2010) Studies on the morphology and spreading of human endothelial cells define key inter- and intramolecular interactions for talin1. <a href="#">Eur J Cell Biol. 89(9):661-73.</a></li> <li>2. Debrand, E. <i>et al.</i> (2012) Mice carrying a complete deletion of the talin2 coding sequence are viable and fertile. <a href="#">Biochem Biophys Res Commun. 426: 190-5.</a></li> <li>3. Manso, A.M. <i>et al.</i> (2013) Talin1 has unique expression versus talin 2 in the heart and modifies the hypertrophic response to pressure overload. <a href="#">J Biol Chem. 288: 4252-64</a></li> <li>4. Liu, J. <i>et al.</i> (2011) Talin1 regulates integrin turnover to promote embryonic epithelial morphogenesis. <a href="#">Mol Cell Biol. 31: 3366-77.</a></li> <li>5. Monkley, S.J. <i>et al.</i> (2011) Endothelial cell talin1 is essential for embryonic angiogenesis. <a href="#">Dev Biol. 349: 494-502.</a></li> <li>6. Ketscher, A. <i>et al.</i> (2014) LSD1 controls metastasis of androgen-independent prostate cancer cells through PXN and LPAR6. <a href="#">Oncogenesis. 3: e120.</a></li> <li>7. Qi, L. <i>et al.</i> (2016) Talin2-mediated traction force drives matrix degradation and cell invasion. <a href="#">J Cell Sci. 129 (19): 3661-74.</a></li> <li>8. Ichikawa, Y. <i>et al.</i> (2017) Modulation of caveolins, integrins and plasma membrane repair proteins in anthracycline-induced heart failure in rabbits. <a href="#">PLoS One. 12 (5): e0177660.</a></li> </ol>
<b>Storage</b>	<p>Store at +4°C or at -20°C if preferred.</p> <p>This product should be stored undiluted.</p> <p>Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
<b>Guarantee</b>	12 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #10040 available at:  
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)  
Goat Anti Mouse IgG (STAR77...) [HRP](#)  
Rabbit Anti Mouse IgG (STAR12...) [RPE](#)  
Rabbit Anti Mouse IgG (STAR8...) [DyLight@800](#)  
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)  
Goat Anti Mouse IgG (STAR76...) [RPE](#)  
Goat Anti Mouse IgG (STAR70...) [FITC](#)  
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)  
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)  
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight@488](#), [DyLight@680](#),  
[DyLight@800](#), [FITC](#), [HRP](#)

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

'M368011:200529'

Printed on 11 Aug 2020

---

© 2020 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)