

Datasheet: MCA47A488

Description:	MOUSE ANTI RAT CD90:Alexa Fluor® 488		
Specificity:	CD90		
Other names:	THY1		
Format:	ALEXA FLUOR® 488		
Product Type:	Monoclonal Antibody		
Clone:	OX-7		
Isotype:	lgG1		
Quantity:	100 TESTS/1ml		

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Rat				
Species Cross	Reacts with: Rabb	it, Mouse, Guinea Pig			
Reactivity	N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications o personal communications from the originators. Please refer to references indicated for further information.				
Product Form	Purified IgG conjugated to Alexa Fluor® 488 - liquid				
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nr	m)	
	Alexa Fluor®488	495	519		
Preparation	Purified IgG prepa supernatant	red by affinity chromatog	raphy on Protein A	from tissue culture	
Buffer Solution	Phosphate buffere	d saline			

Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin				
Approx. Protein Concentrations	IgG concentration 0.05 mg/ml				
Immunogen	Rat Thy1 antigen.				
External Database Links	UniProt: P01830 Related reagents Entrez Gene: 24832 Thy1 Related reagents				
Synonyms	Thy-1				
RRID	AB_324781				
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse NS1 myeloma cell line.				
Specificity	Mouse anti Rat CD90 antibody, clone OX-7 recognizes rat and CD90, also known as Thy1.1, a GPI-anchored membrane protein containing a single V type Ig-like domain CD90 is expressed on a variety of cell types including thymocytes, neuronal cells, stem cells, immature B cells and connective tissues, CD90 is also expressed in T cells in mice.				
	Since Thy1.1 is a monomorphic determinant in rat but polymorphic in mice, clone MRC OX-7 reacts with Thy1.1 mice e.g. AKR and FVB, but not Thy1.2 mice such as CBA and BALB/c. The affinity of the Fab' of MRC OX-7 for rat Thy1 is $3 \times 10^9 \text{m}^{-1}$ and for mouse Thy1.1 is $3 \times 10^8 \text{m}^{-1}$ (1).				
	Mouse anti rat CD90, clone MRC OX-7 has been demonstrated to promote neurite outgrowths on peripherin-stained sympathetic rat neurons, using fluorescence microscopy (<u>Jeng et al. 1998</u>). Clone OX-7 has also been reported to induce glomerular nephritis in Wistar rats (<u>Tamura et al. 1996</u>).				
	This product is routinely tested in flow cytometry on rat thymocytes.				
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.				
References	 Campbell, D.G. <i>et al.</i> (1981) Rat brain Thy-1 glycoprotein. The amino acid sequence, disulphide bonds and an unusual hydrophobic region. <u>Biochem J. 195 (1): 15-30.</u> Bukovský, A. <i>et al.</i> (1983) The localization of Thy-1.1, MRC OX 2 and Ia antigens in the rat ovary and fallopian tube. <u>Immunology. 48 (3): 587-96.</u> 				

399-404.

3. Kawachi, H. et al. (1992) Epitope-specific induction of mesangial lesions with

proteinuria by a MoAb against mesangial cell surface antigen. Clin Exp Immunol. 88 (3):

- 4. Tamura, M. *et al.* (1996) Enhanced glomerular profilin gene and protein expression in experimental mesangial proliferative glomerulonephritis. <u>Biochem Biophys Res Commun.</u> 222 (3): 683-7.
- 5. Banerjee, S.A. *et al.* (1997) An antibody to the tetraspan membrane protein CD9 promotes neurite formation in a partially alpha3beta1 integrin-dependent manner. <u>J. Neurosci.</u> 17 (8): 2756-65.
- 6. Lee, W.S. *et al.* (1998) Thy-1, a novel marker for angiogenesis upregulated by inflammatory cytokines. <u>Circ Res. 82 (8): 845-51.</u>
- 7. Keller, R.K. *et al.* (2004) Formation of 7-dehydrocholesterol-containing membrane rafts *in vitro* and *in vivo*, with relevance to the Smith-Lemli-Opitz syndrome. <u>J Lipid Res. 45:</u> 347-55.
- 8. Rutigliano, J.A. *et al.* (2008) Screening monoclonal antibodies for cross-reactivity in the ferret model of influenza infection. J Immunol Methods. 336: 71-7.
- 9. Stevenson, K.S. *et al.* (2009) Isolation, characterization, and differentiation of thy1.1-sorted pancreatic adult progenitor cell populations. <u>Stem Cells Dev. 18 (10): 1389-98.</u>
- 10. Ohashi, N. *et al.* (2010) Glomerular angiotensinogen is induced in mesangial cells in diabetic rats via reactive oxygen species--ERK/JNK pathways. <u>Hypertens Res.</u> 33:1174-81.
- 11. Biermann, J. *et al.* (2011) Histone deacetylase inhibitors sodium butyrate and valproic acid delay spontaneous cell death in purified rat retinal ganglion cells. <u>Mol Vis. 17:</u> 395-403.
- 12. Freisinger, W. *et al.* (2013) Sensory renal innervation: a kidney-specific firing activity due to a unique expression pattern of voltage-gated sodium channels? <u>Am J Physiol</u> Renal Physiol. 304: F491-7.
- 13. Shimizu T *et al.* (2016) Bioactivity of sol-gel-derived TiO2 coating on polyetheretherketone: *In vitro* and *in vivo* studies. <u>Acta Biomater. 35: 305-17.</u>
- 14. Maia, L. *et al.* (2017) Conditioned medium: a new alternative for cryopreservation of equine umbilical cord mesenchymal stem cells. <u>Cell Biol Int. 41 (3): 239-48.</u>
- 15. Maia, L. *et al.* (2017) A proteomic study of mesenchymal stem cells from equine umbilical cord. Theriogenology. 100: 8-15.
- 16. Zhao, Y. *et al.* (2017) A new electrospun graphene-silk fibroin composite scaffolds for guiding Schwann cells. <u>J Biomater Sci Polym Ed. 28 (18): 2171-85.</u>
- 17. Chang, J.C. *et al.* (2019) Early Immune Response to Acute Gastric Fluid Aspiration in a Rat Model of Lung Transplantation. <u>Exp Clin Transplant</u>. 17 (1): 84-92.
- 18. Huang, X. *et al.* (2019) MRI Tracking of SPIO- and *Fth1*-Labeled Bone Marrow Mesenchymal Stromal Cell Transplantation for Treatment of Stroke. <u>Contrast Media Mol Imaging</u>. 2019: 5184105.
- 19. Kuriyama, T. *et al.* (2020) A novel rat model of inflammatory bowel disease developed using a device created with a 3D printer. <u>Regen Ther. 14: 1-10.</u>
- 20. Cąkała-Jakimowicz, M. & Puzianowska-Kuznicka, M. (2022) Towards Understanding the Lymph Node Response to Skin Infection with Saprophytic *Staphylococcus epidermidis.*. <u>Biomedicines.</u> 10 (5): 1021.
- 21. Huang, S. *et al.* (2022) Hydrogen sulfide supplement preserves mitochondrial function of retinal ganglion cell in a rat glaucoma model. <u>Cell Tissue Res. 389 (2): 171-85.</u>
- 22. Cacciamali, A. *et al.* (2022) Engineered nanoparticles toxicity on adipose tissue derived mesenchymal stem cells: A preliminary investigation Res Vet Sci. 152: 134-49.
- 23. Eweida, A. et al. (2022) Systemically injected bone marrow mononuclear cells

specifically home to axially vascularized tissue engineering constructs. PLoS One. 17 (8): e0272697.

24. Ichinohe, N. et al. (2023) CINC-2 and miR-199a-5p in EVs secreted by transplanted Thy1(+) cells activate hepatocytic progenitor cell growth in rat liver regeneration. Stem Cell Res Ther. 14 (1): 134.

25. Numata-Uematasu, Y. et al. (2023) In vitro myelination using explant culture of dorsal root ganglia: An efficient tool for analyzing peripheral nerve differentiation and disease modeling. PLoS One. 18 (5): e0285897.

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee

12 months from date of despatch

Acknowledgements

This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

Health And Safety Information

Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA47A488 10041

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL: Alexa Fluor® 488 (MCA1209A488)

North & South Tel: +1 800 265 7376

Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21

America

Fax: +1 919 878 3751

Fax: +44 (0)1865 852 739

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M385713:210513'

Printed on 23 May 2025