Datasheet: MCA47FA BATCH NUMBER 1804

| Description: MOUSE ANTI RAT CDS | | |
|---------------------------------|---------------------|--|
| Specificity: | CD90 | |
| Other names: | THY1 | |
| Format: | FITC | |
| Product Type: | Monoclonal Antibody | |
| Clone: | OX-7 | |
| Isotype: | lgG1 | |
| Quantity: | 50 µg | |

Product Details

| Applications | This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> . | | | | | | | |
|-----------------------------|--|----------------------------|--------------------------|--|--|--|--|--|
| | Yes No Not Determined Suggested Dilution | | | | | | | |
| | Flow Cytometry | - | | | | | | |
| | Where this antibody has not been tested for use in a particular technique this does necessarily exclude its use in such procedures. Suggested working dilutions are gi a guide only. It is recommended that the user titrates the antibody for use in their o system using appropriate negative/positive controls. | | | | | | | |
| Target Species | Rat | | | | | | | |
| Species Cross Reactivity | Reacts with: Rabbit, Mouse, Guinea Pig N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. | | | | | | | |
| Product Form | Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid | | | | | | | |
| Max Ex/Em | Fluorophore FITC | Excitation Max (nm) 490 | Emission Max (nm) 525 | | | | | |
| Preparation | Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant | | | | | | | |

| Buffer Solution | Phosphate buffered saline | | |
|-----------------------------------|---|--|--|
| Preservative Stabilisers | 0.09% Sodium Azide 1% Bovine Serum Albumin | | |
| Approx. Protein Concentrations | IgG concentration 0.1 mg/ml | | |
| Immunogen | Rat Thy1 antigen. | | |
| External Database Links | UniProt: P01830 Related reagents Entrez Gene: 24832 Thy1 Related reagents | | |
| Synonyms | Thy-1 | | |
| RRID | AB_566923 | | |
| Fusion Partners | Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS1 myeloma cell line. | | |
| Specificity | Mouse anti Rat CD90 antibody, clone OX-7 recognizes rat and CD90, also known as Thy1.1, a GPI-anchored membrane protein containing a single V type Ig-like domain CD90 is expressed on a variety of cell types including thymocytes, neuronal cells, stem cells, immature B cells and connective tissues, CD90 is also expressed in T cells in mice. | | |
| | Since Thy1.1 is a monomorphic determinant in rat but polymorphic in mice, clone MRC OX-7 reacts with Thy1.1 mice e.g. AKR and FVB, but not Thy1.2 mice such as CBA and BALB/c. The affinity of the Fab' of MRC OX-7 for rat Thy1 is 3×10^9 m- ¹ and for mouse Thy1.1 is 3×10^8 m- ¹ (1). | | |
| | Mouse anti rat CD90, clone MRC OX-7 has been demonstrated to promote neurite outgrowths on peripherin-stained sympathetic rat neurons, using fluorescence microscopy (<u>Jeng <i>et al.</i> 1998</u>). Clone OX-7 has also been reported to induce glomerular nephritis in Wistar rats (<u>Tamura <i>et al.</i> 1996</u>). | | |
| | This product is routinely tested in flow cytometry on rat thymocytes. | | |
| Flow Cytometry | Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul. | | |
| References | Mason, D.W. & Williams, A.F. (1980) The kinetics of antibody binding to membrane antigens in solution and at the cell surface. <u>Biochem J. 187 (1): 1-20.</u> Campbell, D.G. <i>et al.</i> (1981) Rat brain Thy-1 glycoprotein. The amino acid sequence, disulphide bonds and an unusual hydrophobic region. <u>Biochem J. 195 (1): 15-30.</u> Bukovský, A. <i>et al.</i> (1983) The localization of Thy-1.1, MRC OX 2 and Ia antigens in the | | |

rat ovary and fallopian tube. Immunology. 48 (3): 587-96.

4. Lee, W.S. *et al.* (1998) Thy-1, a novel marker for angiogenesis upregulated by inflammatory cytokines. <u>Circ Res. 82 (8): 845-51.</u>

5. Jeng, C.J. *et al.* (1998) Thy-1 is a component common to multiple populations of synaptic vesicles. J Cell Biol. 140 (3): 685-98.

6. Banerjee, S.A. *et al.* (1997) An antibody to the tetraspan membrane protein CD9 promotes neurite formation in a partially alpha3beta1 integrin-dependent manner. J Neurosci. 17 (8): 2756-65.

7. Kawachi, H. *et al.* (1992) Epitope-specific induction of mesangial lesions with proteinuria by a MoAb against mesangial cell surface antigen. <u>Clin Exp Immunol. 88 (3):</u> <u>399-404.</u>

8. Tamura, M. *et al.* (1996) Enhanced glomerular profilin gene and protein expression in experimental mesangial proliferative glomerulonephritis. <u>Biochem Biophys Res Commun.</u> <u>222 (3): 683-7.</u>

9. Stevenson, K.S. *et al.* (2009) Isolation, characterization, and differentiation of thy1.1-sorted pancreatic adult progenitor cell populations. <u>Stem Cells Dev. 18 (10): 1389-98.</u>

10. Biermann, J. *et al.* (2011) Histone deacetylase inhibitors sodium butyrate and valproic acid delay spontaneous cell death in purified rat retinal ganglion cells. <u>Mol Vis. 17:</u> <u>395-403.</u>

11. Keller, R.K. *et al.* (2004) Formation of 7-dehydrocholesterol-containing membrane rafts *in vitro* and *in vivo*, with relevance to the Smith-Lemli-Opitz syndrome. <u>J Lipid Res. 45:</u> <u>347-55.</u>

12. Ohashi, N. *et al.* (2010) Glomerular angiotensinogen is induced in mesangial cells in diabetic rats via reactive oxygen species--ERK/JNK pathways. <u>Hypertens Res.</u> 33:1174-81.

Maia L *et al.* (2016) Conditioned medium: A new alternative for cryopreservation of equine umbilical cord mesenchymal stem cells. <u>Cell Biol Int. Nov 26. [Epub ahead of print]</u>
 Rutigliano, J.A. *et al.* (2008) Screening monoclonal antibodies for cross-reactivity in the ferret model of influenza infection. <u>J Immunol Methods. 336: 71-7.</u>

15. Freisinger, W. *et al.* (2013) Sensory renal innervation: a kidney-specific firing activity due to a unique expression pattern of voltage-gated sodium channels? <u>Am J Physiol</u> <u>Renal Physiol. 304: F491-7.</u>

16. Shimizu T *et al.* (2016) Bioactivity of sol-gel-derived TiO2 coating on polyetheretherketone: *In vitro* and *in vivo* studies. <u>Acta Biomater. 35: 305-17.</u>

17. Maia, L. *et al.* (2017) A proteomic study of mesenchymal stem cells from equine umbilical cord. <u>Theriogenology. 100: 8-15.</u>

18. Chang, J.C. *et al.* (2019) Early Immune Response to Acute Gastric Fluid Aspiration in a Rat Model of Lung Transplantation. <u>Exp Clin Transplant. 17 (1): 84-92.</u>

19. Huang, X. *et al.* (2019) MRI Tracking of SPIO- and *Fth1*-Labeled Bone Marrow Mesenchymal Stromal Cell Transplantation for Treatment of Stroke. <u>Contrast Media Mol</u> <u>Imaging. 2019: 5184105.</u>

20. Zhao, Y. *et al.* (2017) A new electrospun graphene-silk fibroin composite scaffolds for guiding Schwann cells. J Biomater Sci Polym Ed. 28 (18): 2171-85.

Storage Store at +4°C or at -20°C if preferred.

Storage in frost-free freezers is not recommended.

This product should be stored undiluted. This product is photosensitive and should be

| | protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. |
|----------------------------------|---|
| Guarantee | 12 months from date of despatch |
| Health And Safety Information | Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA47FA 10041 |
| Regulatory | For research purposes only |

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:FITC (MCA1209F)

| North & South | Tel: +1 800 265 7376 | Worldwide | Tel: +44 (0)1865 852 700 | Europe | Tel: +49 (0) 89 8090 95 21 |
|---------------|--------------------------------------|-----------|--------------------------------------|--------|--------------------------------------|
| America | Fax: +1 919 878 3751 | | Fax: +44 (0)1865 852 739 | | Fax: +49 (0) 89 8090 95 50 |
| | Email: antibody_sales_us@bio-rad.com | | Email: antibody_sales_uk@bio-rad.com | | Email: antibody_sales_de@bio-rad.com |
| | | | | | |

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M368037:200529'

Printed on 21 Feb 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint