

# Datasheet: MCA47R

Description:	ption: MOUSE ANTI RAT CD90	
Specificity:	CD90	
Other names:	THY1	
Format:	Purified	
<b>Product Type:</b>	Monoclonal Antibody	
Clone:	OX-7	
Isotype:	lgG1	
Quantity:	0.25 mg	

### **Product Details**

## **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			1/50 - 1/100
Immunohistology - Frozen	•			
Immunohistology - Paraffin			•	
ELISA			•	
Immunoprecipitation	-			
Western Blotting	•			
Immunofluorescence	-			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Rat		
Species Cross Reactivity	Reacts with: Rabbit, Mouse, Guinea Pig <b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.		
Product Form	Purified IgG - liquid		
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture		

supernatant

Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	0.09% Sodium Azide		
Carrier Free	Yes		
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml		
Immunogen	Rat Thy1 antigen.		
External Database Links	UniProt: P01830 Related reagents  Entrez Gene: 24832 Thy1 Related reagents		
Synonyms	Thy-1		
RRID	AB_321890		
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse NS1 myeloma cell line.		
Specificity	Mouse anti Rat CD90 antibody, clone OX-7 recognizes rat and CD90, also known as Thy1.1, a GPI-anchored membrane protein containing a single V type Ig-like domain CD90 is expressed on a variety of cell types including thymocytes, neuronal cells, stem cells, immature B cells and connective tissues, CD90 is also expressed in T cells in mice.  Since Thy1.1 is a monomorphic determinant in rat but polymorphic in mice, clone MRC OX-7 reacts with Thy1.1 mice e.g. AKR and FVB, but not Thy1.2 mice such as CBA and BALB/c. The affinity of the Fab' of MRC OX-7 for rat Thy1 is 3 x 10 <sup>9</sup> m- <sup>1</sup> and for mouse Thy1.1 is 3 x 10 <sup>8</sup> m- <sup>1</sup> (1).		
	Mouse anti rat CD90, clone MRC OX-7 has been demonstrated to promote neurite outgrowths on peripherin-stained sympathetic rat neurons, using fluorescence microscopy ( <u>Jeng et al. 1998</u> ). Clone OX-7 has also been reported to induce glomerular nephritis in Wistar rats ( <u>Tamura et al. 1996</u> ).		
Flow Cytometry	This product is routinely tested in flow cytometry on rat thymocytes.  Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.		
References			
1/6161611062	1. Campbell, D.G. <i>et al.</i> (1981) Rat brain Thy-1 glycoprotein. The amino acid sequence, disulphide bonds and an unusual hydrophobic region. <u>Biochem J. 195 (1): 15-30.</u>		

- 2. Bukovský, A. *et al.* (1983) The localization of Thy-1.1, MRC OX 2 and Ia antigens in the rat ovary and fallopian tube. Immunology. 48 (3): 587-96.
- 3. Kawachi, H. *et al.* (1992) Epitope-specific induction of mesangial lesions with proteinuria by a MoAb against mesangial cell surface antigen. <u>Clin Exp Immunol. 88 (3):</u> 399-404.
- 4. Tamura, M. *et al.* (1996) Enhanced glomerular profilin gene and protein expression in experimental mesangial proliferative glomerulonephritis. <u>Biochem Biophys Res Commun.</u> 222 (3): 683-7.
- 5. Banerjee, S.A. *et al.* (1997) An antibody to the tetraspan membrane protein CD9 promotes neurite formation in a partially alpha3beta1 integrin-dependent manner. <u>J Neurosci. 17 (8): 2756-65.</u>
- 6. Lee, W.S. *et al.* (1998) Thy-1, a novel marker for angiogenesis upregulated by inflammatory cytokines. <u>Circ Res. 82 (8): 845-51.</u>
- 7. Keller, R.K. *et al.* (2004) Formation of 7-dehydrocholesterol-containing membrane rafts *in vitro* and *in vivo*, with relevance to the Smith-Lemli-Opitz syndrome. <u>J Lipid Res. 45:</u> 347-55.
- 8. Rutigliano, J.A. *et al.* (2008) Screening monoclonal antibodies for cross-reactivity in the ferret model of influenza infection. <u>J Immunol Methods</u>. 336: 71-7.
- 9. Stevenson, K.S. *et al.* (2009) Isolation, characterization, and differentiation of thy1.1-sorted pancreatic adult progenitor cell populations. <u>Stem Cells Dev. 18 (10): 1389-98.</u>
- 10. Ohashi, N. *et al.* (2010) Glomerular angiotensinogen is induced in mesangial cells in diabetic rats via reactive oxygen species--ERK/JNK pathways. <u>Hypertens Res.</u> 33:1174-81.
- 11. Biermann, J. *et al.* (2011) Histone deacetylase inhibitors sodium butyrate and valproic acid delay spontaneous cell death in purified rat retinal ganglion cells. <u>Mol Vis. 17:</u> 395-403.
- 12. Freisinger, W. *et al.* (2013) Sensory renal innervation: a kidney-specific firing activity due to a unique expression pattern of voltage-gated sodium channels? <u>Am J Physiol</u> Renal Physiol. 304: F491-7.
- 13. Shimizu T *et al.* (2016) Bioactivity of sol-gel-derived TiO2 coating on polyetheretherketone: *In vitro* and *in vivo* studies. <u>Acta Biomater. 35: 305-17.</u>
- 14. Maia, L. *et al.* (2017) Conditioned medium: a new alternative for cryopreservation of equine umbilical cord mesenchymal stem cells. <u>Cell Biol Int. 41 (3): 239-48.</u>
- 15. Maia, L. *et al.* (2017) A proteomic study of mesenchymal stem cells from equine umbilical cord. Theriogenology. 100: 8-15.
- 16. Zhao, Y. *et al.* (2017) A new electrospun graphene-silk fibroin composite scaffolds for guiding Schwann cells. J Biomater Sci Polym Ed. 28 (18): 2171-85.
- 17. Chang, J.C. *et al.* (2019) Early Immune Response to Acute Gastric Fluid Aspiration in a Rat Model of Lung Transplantation. <u>Exp Clin Transplant</u>. 17 (1): 84-92.
- 18. Huang, X. *et al.* (2019) MRI Tracking of SPIO- and *Fth1*-Labeled Bone Marrow Mesenchymal Stromal Cell Transplantation for Treatment of Stroke. <u>Contrast Media Mol Imaging.</u> 2019: 5184105.
- 19. Kuriyama, T. *et al.* (2020) A novel rat model of inflammatory bowel disease developed using a device created with a 3D printer. Regen Ther. 14: 1-10.
- 20. Cąkała-Jakimowicz, M. & Puzianowska-Kuznicka, M. (2022) Towards Understanding the Lymph Node Response to Skin Infection with Saprophytic *Staphylococcus epidermidis.*. <u>Biomedicines</u>. 10 (5): 1021.

- 21. Huang, S. *et al.* (2022) Hydrogen sulfide supplement preserves mitochondrial function of retinal ganglion cell in a rat glaucoma model. <u>Cell Tissue Res. 389 (2): 171-85.</u>
  22. Cacciamali, A. *et al.* (2022) Engineered nanoparticles toxicity on adipose tissue derived mesenchymal stem cells: A preliminary investigation <u>Res Vet Sci. 152: 134-49.</u>
  23. Eweida, A. *et al.* (2022) Systemically injected bone marrow mononuclear cells specifically home to axially vascularized tissue engineering constructs. <u>PLoS One. 17 (8):</u> e0272697.
- 24. Ichinohe, N. *et al.* (2023) CINC-2 and miR-199a-5p in EVs secreted by transplanted Thy1(+) cells activate hepatocytic progenitor cell growth in rat liver regeneration. <u>Stem Cell Res Ther.</u> 14 (1): 134.
- 25. Numata-Uematasu, Y. *et al.* (2023) *In vitro* myelination using explant culture of dorsal root ganglia: An efficient tool for analyzing peripheral nerve differentiation and disease modeling. PLoS One. 18 (5): e0285897.

#### Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA47R">https://www.bio-rad-antibodies.com/SDS/MCA47R</a> 10040
Regulatory	For research purposes only

### Related Products

### **Recommended Secondary Antibodies**

Rabbit Anti Mouse IgG (STAR12...) RPE

Goat Anti Mouse IgG IgA IgM (STAR87...) Alk. Phos., HRP

Goat Anti Mouse IgG (STAR76...) RPE

Rabbit Anti Mouse IgG (STAR13...) <u>HRP</u>

Goat Anti Mouse IgG (STAR70...) FITC

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

Goat Anti Mouse IgG (STAR77...)

Rabbit Anti Mouse IgG (STAR9...)

FITC

**Recommended Negative Controls** 

MOUSE IgG1 NEGATIVE CONTROL (MCA1209)

North & South Tel: +1 800 265 7376 Worldwide Tel: +44 (0)1865 852 700 Tel: +49 (0) 89 8090 95 21 То Europe America Fax: +1 919 878 3751 Fax: +44 (0)1865 852 739 Fax: +49 (0) 89 8090 95 50 find a Email: antibody\_sales\_de@bio-rad.com

Email: antibody\_sales\_us@bio-rad.com Email: antibody\_sales\_uk@bio-rad.com

batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M384239:210513'

### Printed on 21 Feb 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint