

# Datasheet: MCA48GA

**BATCH NUMBER 1610**

<b>Description:</b>	MOUSE ANTI RAT CD8 ALPHA
<b>Specificity:</b>	CD8 ALPHA
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	OX-8
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin	▪			
ELISA			▪	
Immunoprecipitation	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Rat
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant.
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide
<b>Carrier Free</b>	Yes

<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Rat thymocyte membrane glycoproteins.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P07725</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">24930</a>    Cd8a    <a href="#">Related reagents</a></p>
<b>RRID</b>	AB_566918
<b>Fusion Partners</b>	Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS1 myeloma cell line.
<b>Specificity</b>	<p><b>Mouse anti Rat CD8α, clone MRC OX-8</b>, recognizes the rat CD8 alpha cell surface antigen, expressed by a subset of T lymphocytes, most thymocytes and the majority of NK cells.</p> <p>Mouse anti Rat CD8α, clone MRC OX-8 is suitable for use in <i>in vitro</i> blocking studies (<a href="#">Popov et al.2001</a>).</p> <p>Mouse anti Rat CD8α, clone MRC OX-8 has been described reacting with paraffin-embedded material following PLP Fixation (periodate-lysine paraformaldehyde) (<a href="#">Whiteland et al. 1995</a>).</p> <p>Mouse anti Rat CD8α, clone MRC OX-8 is routinely tested in flow cytometry on rat splenocytes.</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
<b>Immunohistology</b>	<p>This product does not require protein digestion pre-treatment of paraffin embedded sections.</p> <p>This product does not require antigen retrieval using heat treatment prior to staining of paraffin embedded sections.</p>
<b>References</b>	<ol style="list-style-type: none"> <li>1. Brideau, R.J. <i>et al.</i> (1980) Two subsets of rat T lymphocytes defined with monoclonal antibodies. <a href="#">Eur J Immunol. 10 (8): 609-15.</a></li> <li>2. Lyscom, N. &amp; Brueton, M.J. (1982) Intraepithelial, lamina propria and Peyer's patch lymphocytes of the rat small intestine: isolation and characterization in terms of immunoglobulin markers and receptors for monoclonal antibodies. <a href="#">Immunology. 45 (4): 775-83.</a></li> <li>3. Whiteland, J.L. <i>et al.</i> (1995) Immunohistochemical detection of T-cell subsets and other leukocytes in paraffin-embedded rat and mouse tissues with monoclonal antibodies. <a href="#">J Histochem Cytochem. 43 (3): 313-20.</a></li> <li>4. Bukovský A <i>et al.</i> (1984) Association of some cell surface antigens of lymphoid cells</li> </ol>

- and cell surface differentiation antigens with early rat pregnancy. [Immunology. 52 \(4\): 631-40.](#)
5. Popov, I. *et al.* (2001) The effect of an anti-HLA-B27 immune response on CTL recognition of Chlamydia. [J Immunol. 167 \(6\): 3375-82.](#)
  6. Mitnacht, R. *et al.* (1998) Opposite CD4/CD8 lineage decisions of CD4+8+ mouse and rat thymocytes to equivalent triggering signals: correlation with thymic expression of a truncated CD8 alpha chain in mice but not rats. [J Immunol. 160 \(2\): 700-7.](#)
  7. Thomas, M.L. & Green, J.R. (1983) Molecular nature of the W3/25 and MRC OX-8 marker antigens for rat T lymphocytes: comparisons with mouse and human antigens. [Eur J Immunol. 13 \(10\): 855-8.](#)
  8. Torres-Nagel, N. *et al.* (1992) Differential thymus dependence of rat CD8 isoform expression. [Eur J Immunol. 22 \(11\): 2841-8.](#)
  9. Zilka, N. *et al.* (2009) Human misfolded truncated tau protein promotes activation of microglia and leukocyte infiltration in the transgenic rat model of tauopathy. [J Neuroimmunol. 209 \(1-2\): 16-25.](#)
  10. Schwartzkopff, J. *et al.* (2010) NK cell depletion delays corneal allograft rejection in baby rats. [Mol Vis. 16: 1928-35.](#)
  11. Banerjee, S. *et al.* (2003) Development of organised conjunctival leucocyte aggregates after corneal transplantation in rats. [Br J Ophthalmol. 87: 1515-22.](#)
  12. Sanchez-Guajardo, V. (2010) Microglia acquire distinct activation profiles depending on the degree of alpha-synuclein neuropathology in a rAAV based model of Parkinson's disease. [PLoS One. 5: e8784.](#)
  13. Maenz, M. *et al.* (2011) A comprehensive flow-cytometric analysis of graft infiltrating lymphocytes, draining lymph nodes and serum during the rejection phase in a fully allogeneic rat cornea transplant model. [Mol Vis. 2011 Feb 8;17:420-9.](#)
  14. Granados-Durán P *et al.* (2015) Neuroinflammation induced by intracerebroventricular injection of microbial neuraminidase. [Front Med \(Lausanne\). 2: 14.](#)
  15. Nuccitelli R *et al.* (2015) Nanoelectroablation of Murine Tumors Triggers a CD8-Dependent Inhibition of Secondary Tumor Growth. [PLoS One. 10 \(7\): e0134364.](#)
  16. Zhang, Z.M. *et al.* (2016) Lesional accumulation of CD8(+) cells in sciatic nerves of experimental autoimmune neuritis rats. [Neurol Sci. 37 \(2\): 199-203.](#)
  17. Pamukcu, O. *et al.* (2016) Anti-inflammatory role of obestatin in autoimmune myocarditis. [Clin Exp Pharmacol Physiol. 43 \(1\): 47-55.](#)
  18. Dabrowska, S. *et al.* (2019) Human bone marrow mesenchymal stem cell-derived extracellular vesicles attenuate neuroinflammation evoked by focal brain injury in rats. [J Neuroinflammation. 16 \(1\): 216.](#)
  19. Arndt, T. *et al.* (2013) A variable CD3<sup>+</sup> T-cell frequency in peripheral blood lymphocytes associated with type 1 diabetes mellitus development in the LEW.1AR1-iddm rat. [PLoS One. 8 \(5\): e64305.](#)
  20. James, R.E. *et al.* (2020) Persistent elevation of intrathecal pro-inflammatory cytokines leads to multiple sclerosis-like cortical demyelination and neurodegeneration. [Acta Neuropathol Commun. 8 \(1\): 66.](#)

---

#### Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

<b>Guarantee</b>	12 months from date of despatch
<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA48GA">https://www.bio-rad-antibodies.com/SDS/MCA48GA</a> 10040
<b>Regulatory</b>	For research purposes only

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	<a href="#">RPE</a>
Goat Anti Mouse IgG IgA IgM (STAR87...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (STAR76...)	<a href="#">RPE</a>
Rabbit Anti Mouse IgG (STAR13...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (STAR70...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (H/L) (STAR117...)	<a href="#">Alk. Phos.</a> , <a href="#">DyLight®488</a> , <a href="#">DyLight®550</a> , <a href="#">DyLight®650</a> , <a href="#">DyLight®680</a> , <a href="#">DyLight®800</a> , <a href="#">FITC</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR9...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (STAR77...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (Fc) (STAR120...)	<a href="#">FITC</a> , <a href="#">HRP</a>

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA1209\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)  
'M368059:200529'

Printed on 05 Feb 2024