

Datasheet: MCA48R

Description:	MOUSE ANTI RAT CD8 ALPHA
Specificity:	CD8 ALPHA
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	OX-8
Isotype:	IgG1
Quantity:	0.25 mg

Product Details

RRID AB_2260088

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin	▪			
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting	▪			
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species Rat

Product Form Purified IgG - liquid

Preparation Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant.

Buffer Solution Phosphate buffered saline

Preservative Stabilisers 0.09% Sodium Azide

Carrier Free Yes

Approx. Protein Concentrations IgG concentration 1.0 mg/ml

Immunogen Rat thymocyte membrane glycoproteins.

External Database Links	UniProt: P07725 Related reagents Entrez Gene: 24930 Cd8a Related reagents
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS1 myeloma cell line.
Specificity	<p>Mouse anti Rat CD8α, clone MRC OX-8, recognizes the rat CD8 alpha cell surface antigen, expressed by a subset of T lymphocytes, most thymocytes and the majority of NK cells.</p> <p>Clone MRC OX-8 is suitable for use in <i>in vitro</i> blocking studies (Popov et al.2001).</p> <p>Mouse anti Rat CD8α, clone MRC OX-8 has been described reacting with paraffin-embedded material following PLP Fixation (periodate-lysine paraformaldehyde) (Whiteland et al. 1995).</p> <p>Mouse anti Rat CD8α, clone MRC OX-8 is routinely tested in flow cytometry on rat splenocytes.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
Immunohistology	This product does not require protein digestion pre-treatment of paraffin embedded sections. This product does not require antigen retrieval using heat treatment prior to staining of paraffin embedded sections.
References	<ol style="list-style-type: none"> 1. Brideau, R.J. <i>et al.</i> (1980) Two subsets of rat T lymphocytes defined with monoclonal antibodies. Eur J Immunol. 10 (8): 609-15. 2. Lyscom, N. & Brueton, M.J. (1982) Intraepithelial, lamina propria and Peyer's patch lymphocytes of the rat small intestine: isolation and characterization in terms of immunoglobulin markers and receptors for monoclonal antibodies. Immunology. 45 (4): 775-83. 3. Whiteland, J.L. <i>et al.</i> (1995) Immunohistochemical detection of T-cell subsets and other leukocytes in paraffin-embedded rat and mouse tissues with monoclonal antibodies. J Histochem Cytochem. 43 (3): 313-20. 4. Bukovský A <i>et al.</i> (1984) Association of some cell surface antigens of lymphoid cells and cell surface differentiation antigens with early rat pregnancy. Immunology. 52 (4): 631-40. 5. Popov, I. <i>et al.</i> (2001) The effect of an anti-HLA-B27 immune response on CTL recognition of Chlamydia. J Immunol. 167 (6): 3375-82. 6. Mitnacht, R. <i>et al.</i> (1998) Opposite CD4/CD8 lineage decisions of CD4+8+ mouse and rat thymocytes to equivalent triggering signals: correlation with thymic expression of a truncated CD8 alpha chain in mice but not rats. J Immunol. 160 (2): 700-7. 7. Thomas, M.L. & Green, J.R. (1983) Molecular nature of the W3/25 and MRC OX-8 marker antigens for rat T lymphocytes: comparisons with mouse and human antigens. Eur J Immunol. 13 (10): 855-8. 8. Torres-Nagel, N. <i>et al.</i> (1992) Differential thymus dependence of rat CD8 isoform expression. Eur J Immunol. 22 (11): 2841-8. 9. Zilka, N. <i>et al.</i> (2009) Human misfolded truncated tau protein promotes activation of microglia and leukocyte infiltration in the transgenic rat model of tauopathy. J Neuroimmunol. 209 (1-2): 16-25. 10. Schwartzkopff, J. <i>et al.</i> (2010) NK cell depletion delays corneal allograft rejection in baby rats. Mol Vis. 16: 1928-35. 11. Banerjee, S. <i>et al.</i> (2003) Development of organised conjunctival leucocyte aggregates after

- corneal transplantation in rats. [Br J Ophthalmol. 87: 1515-22.](#)
12. Sanchez-Guajardo, V. (2010) Microglia acquire distinct activation profiles depending on the degree of alpha-synuclein neuropathology in a rAAV based model of Parkinson's disease. [PLoS One. 5: e8784.](#)
13. Maenz, M. *et al.* (2011) A comprehensive flow-cytometric analysis of graft infiltrating lymphocytes, draining lymph nodes and serum during the rejection phase in a fully allogeneic rat cornea transplant model. [Mol Vis. 2011 Feb 8;17:420-9.](#)
14. Granados-Durán P *et al.* (2015) Neuroinflammation induced by intracerebroventricular injection of microbial neuraminidase. [Front Med \(Lausanne\). 2: 14.](#)
15. Nuccitelli R *et al.* (2015) Nanoelectroablation of Murine Tumors Triggers a CD8-Dependent Inhibition of Secondary Tumor Growth. [PLoS One. 10 \(7\): e0134364.](#)
16. Zhang, Z.M. *et al.* (2016) Lesional accumulation of CD8(+) cells in sciatic nerves of experimental autoimmune neuritis rats. [Neurol Sci. 37 \(2\): 199-203.](#)
17. Pamukcu, O. *et al.* (2016) Anti-inflammatory role of obestatin in autoimmune myocarditis. [Clin Exp Pharmacol Physiol. 43 \(1\): 47-55.](#)

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 18 months from date of despatch.

Health And Safety Information Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

- Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)
- Goat Anti Mouse IgG (STAR77...) [HRP](#)
- Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
- Rabbit Anti Mouse IgG (STAR8...) [DyLight®800](#)
- Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
- Goat Anti Mouse IgG (STAR76...) [RPE](#)
- Goat Anti Mouse IgG (STAR70...) [FITC](#)
- Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
- Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
- Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®680](#), [DyLight®800](#), [FITC](#), [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA1209\)](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50

Printed on 20 May 2019
