

Datasheet: MCA500A647

Description:	RAT ANTI MOUSE CD3:Alexa Fluor® 647
Specificity:	CD3
Format:	ALEXA FLUOR® 647
Product Type:	Monoclonal Antibody
Clone:	KT3
Isotype:	IgG2a
Quantity:	100 TESTS/1ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse						
Product Form	Purified IgG conjugated to Alexa Fluor® 647 - liquid						
Max Ex/Em	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>Alexa Fluor®647</td> <td>650</td> <td>665</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	Alexa Fluor®647	650	665
Fluorophore	Excitation Max (nm)	Emission Max (nm)					
Alexa Fluor®647	650	665					
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant						
Buffer Solution	Phosphate buffered saline						
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin						
Approx. Protein Concentrations	IgG concentration 0.05 mg/ml						
Immunogen	CBAT6 thymocytes						

External Database**Links****UniProt:**

[P22646](#) [Related reagents](#)
[P24161](#) [Related reagents](#)
[P11942](#) [Related reagents](#)
[P29020](#) [Related reagents](#)
[P04235](#) [Related reagents](#)

Entrez Gene:

[12501](#) Cd3e [Related reagents](#)
[12503](#) Cd247 [Related reagents](#)
[12500](#) Cd3d [Related reagents](#)
[12502](#) Cd3g [Related reagents](#)
[12503](#) Cd247 [Related reagents](#)

Synonyms

Cd3z, T3d, Tcrz

RRID

AB_324732

Fusion Partners

Spleen cells from immunized SD rats were fused with cells of the NS0 mouse myeloma cell line.

Specificity

Rat anti Mouse CD3 antibody, clone KT3 recognizes the mouse CD3 antigen, expressed by mature T cells. Rat anti Mouse CD3 antibody, clone KT3 may be used to trigger proliferation and cytotoxicity of CD3 positive cells ([Tomonari 1988](#)).

NB. For optimal staining incubations should be performed at room temperature.

Flow Cytometry

Use 10ul of the suggested working dilution to label 10^6 cells or 100ul whole blood.

The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR ([BUF041A/B](#))

References

1. Tomonari, K. (1988) A rat antibody against a structure functionally related to the mouse T-cell receptor/T3 complex. [Immunogenetics. 28 \(6\): 455-8.](#)
2. Lacroix-Lamandé, S. *et al.* (2002) Role of gamma interferon in chemokine expression in the ileum of mice and in a murine intestinal epithelial cell line after *Cryptosporidium parvum* infection. [Infect Immun. 70 \(4\): 2090-9.](#)
3. Tomonari, K. & Lovering, E. (1988) T-cell receptor-specific monoclonal antibodies against a V beta 11-positive mouse T-cell clone. [Immunogenetics. 28 \(6\): 445-51.](#)
4. Haroon, F. *et al.*, (2011) Gp130-dependent astrocytic survival is critical for the control of autoimmune central nervous system inflammation. [J Immunol. 186: 6521-31.](#)
5. Kumar, L. *et al.* (2002) Differential role of SLP-76 domains in T cell development and function. [Proc Natl Acad Sci U S A. 99: 884-9.](#)
6. Hirsh, M. *et al.* (2004) Response of lung gammadelta T cells to experimental sepsis in mice. [Immunology. 112: 153-60.](#)
7. Tarlton, J.F. *et al.* (2000) The role of up-regulated serine proteases and matrix

- metalloproteinases in the pathogenesis of a murine model of colitis. [Am J Pathol. 157: 1927-35.](#)
8. Lazarovits, A.I. *et al.* (1999) Mechanisms of induction of renal allograft tolerance in CD45RB-treated mice. [Kidney Int. 55: 1303-10.](#)
9. Bauer, D. *et al.* (2009) Amniotic membrane transplantation induces apoptosis in T lymphocytes in murine corneas with experimental herpetic stromal keratitis [Invest Ophthalmol Vis Sci. 50: 3188-98.](#)
10. Heitmann, S. *et al.* (1999) Immunohistological characterization of leukocytes in the lungs of healthy mice and after bacterial intratracheal infection. [Lab Anim. 33: 288-94.](#)
11. Hare, K.J. *et al.* (2003) Modeling TCR signaling complex formation in positive selection. [J Immunol. 171: 2825-31.](#)
12. Erlandsson, L. *et al.* (2004) Impaired B-1 and B-2 B cell development and atypical splenic B cell structures in IL-7 receptor-deficient mice. [Eur J Immunol. 34: 3595-603.](#)
13. Severinová, J. *et al.* (2005) Co-inoculation of *Borrelia afzelii* with tick salivary gland extract influences distribution of immunocompetent cells in the skin and lymph nodes of mice. [Folia Microbiol \(Praha\). 50: 457-63.](#)
14. Rothhammer, V. *et al.* (2011) Th17 lymphocytes traffic to the central nervous system independently of $\alpha 4$ integrin expression during EAE. [J Exp Med. 208 \(12\): 2465-76.](#)
15. Salem, M. *et al.* (2011) Interferon regulatory factor-7 modulates experimental autoimmune encephalomyelitis in mice. [J Neuroinflammation. 8: 181.](#)
16. Hoeksema, M.A. *et al.* (2014) Targeting macrophage Histone deacetylase 3 stabilizes atherosclerotic lesions. [EMBO Mol Med. pii: e201404170.](#)
17. Scheinert, R.B. *et al.* (2016) Therapeutic effects of stress-programmed lymphocytes transferred to chronically stressed mice. [Prog Neuropsychopharmacol Biol Psychiatry. Apr 21. pii: S0278-5846\(16\)30056-2. \[Epub ahead of print\]](#)
18. Schaut, R.G. *et al.* (2015) Bovine viral diarrhea virus type 2 *in vivo* infection modulates TLR4 responsiveness in differentiated myeloid cells which is associated with decreased MyD88 expression. [Virus Res. 208: 44-55.](#)
19. Janssen, E. *et al.* (2016) A DOCK8-WIP-WASp complex links T cell receptors to the actin cytoskeleton. [J Clin Invest. 126 \(10\): 3837-51.](#)
20. Teeling, J.L. *et al.* (2012) Intracerebral immune complex formation induces inflammation in the brain that depends on Fc receptor interaction. [Acta Neuropathol. 124 \(4\): 479-90.](#)
21. Van Aelst, L.N. *et al.* (2016) RNA Profiling in Human and Murine Transplanted Hearts: Identification and Validation of Therapeutic Targets for Acute Cardiac and Renal Allograft Rejection. [Am J Transplant. 16 \(1\): 99-110.](#)
22. Kim, I. *et al.* (2016) Immunological characterization of de novo and recall alloantibody suppression by CTLA4Ig in a mouse model of allosensitization. [Transpl Immunol. 38: 84-92.](#)
23. Massa, M.G. *et al.* (2017) Testosterone Differentially Affects T Cells and Neurons in Murine and Human Models of Neuroinflammation and Neurodegeneration. [Am J Pathol. 187 \(7\): 1613-22.](#)
24. Zamudio, F. *et al.* (2020) TDP-43 mediated blood-brain barrier permeability and leukocyte infiltration promote neurodegeneration in a low-grade systemic inflammation mouse model. [J Neuroinflammation. 17 \(1\): 283.](#)
25. Yun, M. *et al.* (2020) Enriched-Baicalein Attenuates Allergy in Cells and Mice [Ev-Based Comp Alt Med.. 2020: 1-8.](#)

26. Granadillo, M. *et al.* (2019) Impact on antitumor response using a new adjuvant preparation as a component of a human papillomavirus type 16 therapeutic vaccine candidate. [Vaccine. 37 \(30\): 3957-60.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee 12 months from date of despatch

Acknowledgements This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchased product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

Health And Safety Information Material Safety Datasheet documentation #10041 available at: 10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL:Alexa Fluor® 647 \(MCA6005A647\)](#)

Recommended Useful Reagents

[MOUSE SEROBLOCK FcR \(BUF041A\)](#)

[MOUSE SEROBLOCK FcR \(BUF041B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M385742:210513'

Printed on 15 Oct 2021