

Datasheet: MCA503

BATCH NUMBER 162950

Description:	RAT ANTI HUMAN CD18
Specificity:	CD18
Other names:	INTEGRIN BETA 2 CHAIN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	YFC118.3
Isotype:	IgG2b
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen (1)	▪			1/1000 - 1/5000
Immunohistology - Paraffin		▪		
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.

Target Species	Human
Species Cross Reactivity	<p>Reacts with: Dog, Guinea Pig</p> <p>N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for</p>

further information.

Product Form	Purified IgG - liquid
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1 mg/ml
Immunogen	Human neutrophils
External Database Links	UniProt: P05107 Related reagents Entrez Gene: 3689 ITGB2 Related reagents
Synonyms	CD18, MFI7
RRID	AB_321325
Fusion Partners	Spleen cells from immunized LOU rats were fused with cells of the rat Y3/Ag.1.2.3 myeloma cell line
Specificity	<p>Rat anti Human CD18 antibody, clone YFC118.3 was clustered at the Fourth International Workshop on Leucocyte Differentiation Antigens (code number N221) as recognizing the CD18 antigen.</p> <p>CD18 is an integral membrane glycoprotein of ~95 kDa, also known as the beta 2 chain, of the LFA-1 complex. CD18 links non-covalently to either CD11a, b or c molecules forming the heteromeric LFA-1 complex. CD18 acts as the receptor for ICAM-1 and is important for cell adhesion and cell-cell interactions (Reina & Espel 2017).</p> <p>Rat anti Human CD18 antibody, clone YFC118.3 demonstrates strong reactivity with leucocytes and is not reactive with platelets.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
Histology Positive Control Tissue	Human Tonsil
References	1. Bindon, C.I. <i>et al.</i> (1988) Importance of antigen specificity for complement-mediated lysis by monoclonal antibodies. Eur J Immunol. 18 (10): 1507-14.

2. Chabanne, L. *et al.* (1994) Screening of 78 monoclonal antibodies directed against human leukocyte antigens for cross-reactivity with surface markers on canine lymphocytes. [Tissue Antigens. 43 \(3\): 202-5.](#)
3. Sampaio, W.M. *et al.* (2007) *In vitro* binding and survival assays of Leishmania parasites to peripheral blood monocytes and monocyte-derived macrophages isolated from dogs naturally and experimentally infected with *Leishmania chagasi*. [BMC Vet Res. 3:11.](#)
4. Canalli, A.A. *et al.* (2011) Participation of Mac-1, LFA-1 and VLA-4 integrins in the *in vitro* adhesion of sickle cell disease neutrophils to endothelial layers, and reversal of adhesion by simvastatin. [Haematologica. 96: 526-33.](#)
5. Gonçalves, R. *et al.* (2005) A sensitive flow cytometric methodology for studying the binding of *L. chagasi* to canine peritoneal macrophages. [BMC Infect Dis. 5:39.](#)
6. Zimmerman, K.L. *et al.* (2013) Leukocyte adhesion deficiency type I in a mixed-breed dog. [J Vet Diagn Invest. 25: 291-6.](#)
7. Araújo, M.S. *et al.* (2011) Immunological changes in canine peripheral blood leukocytes triggered by immunization with first or second generation vaccines against canine visceral leishmaniasis. [Vet Immunol Immunopathol. 141: 64-75.](#)
8. Garland, R.J. *et al.* (2002) Human CD8+ CTL recognition and *in vitro* lysis of herpes simplex virus-infected cells by a non-MHC restricted mechanism. [Scand J Immunol. 55: 61-9.](#)
9. Grote, K. *et al.* (2007) The angiogenic factor CCN1 promotes adhesion and migration of circulating CD34+ progenitor cells: potential role in angiogenesis and endothelial regeneration. [Blood. 110: 877-85.](#)
10. Holst, B.S. *et al.* (2010) Expression of four canine leukocyte adhesion factors in fresh and stored whole blood samples evaluated using a no-lyse, no-wash method. [Vet Immunol Immunopathol. 139: 271-6.](#)
11. Kupatt, C. *et al.* (1999) Tumor necrosis factor-alpha contributes to ischemia- and reperfusion-induced endothelial activation in isolated hearts. [Circ Res. 84: 392-400.](#)
12. Marconato, L. (2013) The dog as a possible animal model for human non-Hodgkin lymphoma: a review. [Hematol Oncol. 31: 1-9.](#)
13. Spring, F.A. *et al.* (2001) Intercellular adhesion molecule-4 binds alpha(4)beta(1) and alpha(V)-family integrins through novel integrin-binding mechanisms. [Blood. 98: 458-66.](#)
14. Salvatierra, A. *et al.* (2001) Antithrombin III prevents early pulmonary dysfunction after lung transplantation in the dog. [Circulation. 104: 2975-80.](#)
15. Mizuno, T. *et al.* (1997) cDNA cloning and chromosomal localization of the human telencephalin and its distinctive interaction with lymphocyte function-associated antigen-1. [J Biol Chem. 272: 1156-63.](#)
16. Lana, S. *et al.* (2006) Diagnosis of mediastinal masses in dogs by flow cytometry. [J Vet Intern Med. 20: 1161-5.](#)
17. Waché, Y.J. *et al.* (2009) The mycotoxin deoxynivalenol inhibits the cell surface expression of activation markers in human macrophages. [Toxicology. 262: 239-44.](#)
18. Crosby, H.A. *et al.* (2009) Adhesion of human haematopoietic (CD34+) stem cells to human liver compartments is integrin and CD44 dependent and modulated by CXCR3 and CXCR4. [J Hepatol. 51: 734-49.](#)
19. Levy, O. *et al.* (2015) Apolipoprotein E promotes subretinal mononuclear phagocyte survival and chronic inflammation in age-related macular degeneration. [EMBO Mol Med. pii: e201404524.](#)

20. Alex, J. *et al.* (2005) Pretreatment with hyperbaric oxygen and its effect on neuropsychometric dysfunction and systemic inflammatory response after cardiopulmonary bypass: a prospective randomized double-blind trial. [J Thorac Cardiovasc Surg. 130: 1623-30.](#)
21. Sutcliffe, J.E.S. *et al.* (2017) Changes in the extracellular matrix surrounding human chronic wounds revealed by 2-photon imaging. [Int Wound J. Jul 20 \[Epub ahead of print\].](#)
22. Cremer, S.E. *et al.* (2019) Proteomic profiling of the thrombin-activated canine platelet secretome (CAPS). [PLoS One. 14 \(11\): e0224891.](#)
23. Wolf-Ringwall, A. *et al.* (2020) Prospective evaluation of flow cytometric characteristics, histopathologic diagnosis and clinical outcome in dogs with naïve B-cell lymphoma treated with a 19-week CHOP protocol. [Vet Comp Oncol. 18 \(3\): 342-52.](#)

Storage	<p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.</p>
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA503
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	DyLight®800
Rabbit Anti Rat IgG (STAR17...)	FITC
Rabbit Anti Rat IgG (STAR21...)	HRP
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®550 , DyLight®650 , DyLight®800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR73...)	RPE
Goat Anti Rat IgG (STAR72...)	HRP

Recommended Negative Controls

[RAT IgG2b NEGATIVE CONTROL \(MCA6006GA\)](#)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M384267:210513'

Printed on 15 Aug 2025

