

Datasheet: MCA506PET

Description:	RAT ANTI HUMAN CD235a:RPE
Specificity:	CD235a
Other names:	GLYCOPHORIN A
Format:	RPE
<b>Product Type:</b>	Monoclonal Antibody
Clone:	YTH89.1
Isotype:	lgG2b
Quantity:	25 TESTS

### **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	-			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human			
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized			
Reconstitution	Reconstitute in 0.25 ml disilled water			
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)	
	RPE 488nm laser	496	578	
Buffer Solution	Phosphate buffered s	aline		
Preservative	0.09% Sodium Azide			
Stabilisers	1% Bovine Serum Albumin			
	5% Sucrose			
External Database				
Links	UniProt:			

## P02724 Related reagents

## **Entrez Gene:**

2993 GYPA Related reagents

Synonyms	GPA
RRID	AB_1101586
Fusion Partners	Spleen cells from immunized DA rats were fused with cells of the Y3/Ag.1.2.3.
Specificity	Rat anti Human CD235a antibody, clone YTH89.1 recognizes glycophorin A, a major sialoglycoprotein of the human erythrocyte membrane.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
References	1. Outram, S. <i>et al.</i> (1988) Erythromyeloid lineage fidelity is conserved in erythroleukaemia. Leuk Res. 12 (8): 651-7.  2. Jokiranta, T.S. & Meri, S. (1993) Biotinylation of monoclonal antibodies prevents their ability to activate the classical pathway of complement. J Immunol. 151 (4): 2124-31.  3. Basu, S. (2010) Erythrocyte membrane defects and asymmetry in paroxysmal nocturnal hemoglobinuria and myelodysplastic syndrome. Hematology. 15: 236-9.  4. Hoang, T. <i>et al.</i> (1996) Opposing effects of the basic helix-loop-helix transcription factor SCL on erythroid and monocytic differentiation. Blood. 87: 102-11.  5. Lahlil, R. <i>et al.</i> (2004) SCL assembles a multifactorial complex that determines glycophorin A expression. Mol Cell Biol. 24: 1439-52.  6. Tiziani, S. <i>et al.</i> (2009) Metabolomic profiling of drug responses in acute myeloid leukaemia cell lines. PLoS One. 2009;4(1):e4251.  7. Saha, S. <i>et al.</i> (2011) Elevated levels of redox regulators, membrane-bound globin chains, and cytoskeletal protein fragments in hereditary spherocytosis erythrocyte proteome. Eur J Haematol. 87: 259-66.  8. Challier, J.C. <i>et al.</i> (2005) Immunocytological evidence for hematopoiesis in the early human placenta. Placenta. 26: 282-8.  9. Huang, Y.C. <i>et al.</i> (2009) Oral small-molecule tyrosine kinase inhibitor midostaurin (PKC412) inhibits growth and induces megakaryocytic differentiation in human leukemia cells. Toxicol In Vitro. 23: 979-85.  10. Lucky AB <i>et al.</i> (2016) Plasmodium knowlesi Skeleton-Binding Protein 1 Localizes to the 'Sinton and Mulligan' Stipplings in the Cytoplasm of Monkey and Human Erythrocytes. PLoS One. 11 (10): e0164272.  11. Babiker, A.A. <i>et al.</i> (2002) Transfer of prostasomal CD59 to CD59-deficient red blood cells results in protection against complement-mediated hemolysis. Am J Reprod Immunol. 47 (3): 183-92.
Storage	Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.

This product should be stored undiluted.

DO NOT FREEZE. This product is photosensitive and should be protected from light.

Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of reconstitution.
Health And Safety Information	Material Safety Datasheet documentation #10075 available at: 10075: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf</a>
Regulatory	For research purposes only

## **Related Products**

# **Recommended Negative Controls**

RAT IgG2b NEGATIVE CONTROL:RPE (MCA6006PE)

### **Recommended Useful Reagents**

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

From March 15, 2021, we will no longer supply printed datasheets with our products. Look out for updates on how to access your digital version at bio-rad-antibodies.com 'M337097:181214'

#### Printed on 10 Feb 2021

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint