

## Datasheet: MCA506SBB700

**BATCH NUMBER 100008104**

<b>Description:</b>	RAT ANTI HUMAN CD235a:StarBright Blue 700
<b>Specificity:</b>	CD235a
<b>Other names:</b>	GLYCOPHORIN A
<b>Format:</b>	StarBright Blue 700
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	YTH89.1
<b>Isotype:</b>	IgG2b
<b>Quantity:</b>	100 TESTS/0.5ml

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Human		
<b>Product Form</b>	Purified IgG conjugated to StarBright Blue 700 - liquid		
<b>Max Ex/Em</b>	<b>Fluorophore</b>	<b>Excitation Max (nm)</b>	<b>Emission Max (nm)</b>
	StarBright Blue 700	473	703
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
<b>Buffer Solution</b>	Phosphate buffered saline		
<b>Preservative</b>	0.09% Sodium Azide (NaN <sub>3</sub> )		
<b>Stabilisers</b>	1% Bovine Serum Albumin		
	0.1% Pluronic F68		
	0.1% PEG 3350		

---

**External Database****Links****UniProt:**[P02724](#)[Related reagents](#)**Entrez Gene:**[2993](#)

GYPA

[Related reagents](#)

---

**Synonyms**

GPA

---

**Fusion Partners**

Spleen cells from immunized DA rats were fused with cells of the Y3/Ag.1.2.3.

---

**Specificity****Rat anti Human CD235a antibody, clone YTH89.1** recognizes glycophorin A, a major sialoglycoprotein of the human erythrocyte membrane.

---

**Flow Cytometry**Use 5µl of the suggested working dilution to label 10<sup>6</sup> cells in 100µl. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.

---

**References**

1. Outram, S. *et al.* (1988) Erythromyeloid lineage fidelity is conserved in erythroleukaemia. [Leuk Res. 12 \(8\): 651-7.](#)
2. Jokiranta, T.S. & Meri, S. (1993) Biotinylation of monoclonal antibodies prevents their ability to activate the classical pathway of complement. [J Immunol. 151 \(4\): 2124-31.](#)
3. Hoang, T. *et al.* (1996) Opposing effects of the basic helix-loop-helix transcription factor SCL on erythroid and monocytic differentiation. [Blood. 87: 102-11.](#)
4. Babiker, A.A. *et al.* (2002) Transfer of prostasomal CD59 to CD59-deficient red blood cells results in protection against complement-mediated hemolysis. [Am J Reprod Immunol. 47 \(3\): 183-92.](#)
5. Lahlil, R. *et al.* (2004) SCL assembles a multifactorial complex that determines glycophorin A expression. [Mol Cell Biol. 24: 1439-52.](#)
6. Challier, J.C. *et al.* (2005) Immunocytological evidence for hematopoiesis in the early human placenta. [Placenta. 26: 282-8.](#)
7. Huang, Y.C. *et al.* (2009) Oral small-molecule tyrosine kinase inhibitor midostaurin (PKC412) inhibits growth and induces megakaryocytic differentiation in human leukemia cells. [Toxicol In Vitro. 23: 979-85.](#)
8. Tiziani, S. *et al.* (2009) Metabolomic profiling of drug responses in acute myeloid leukaemia cell lines. [PLoS One. 2009;4\(1\):e4251.](#)
9. Basu, S. (2010) Erythrocyte membrane defects and asymmetry in paroxysmal nocturnal hemoglobinuria and myelodysplastic syndrome. [Hematology. 15: 236-9.](#)
10. Saha, S. *et al.* (2011) Elevated levels of redox regulators, membrane-bound globin chains, and cytoskeletal protein fragments in hereditary spherocytosis erythrocyte proteome. [Eur J Haematol. 87: 259-66.](#)
11. Lucky AB *et al.* (2016) Plasmodium knowlesi Skeleton-Binding Protein 1 Localizes to the 'Sinton and Mulligan' Stipplings in the Cytoplasm of Monkey and Human Erythrocytes. [PLoS One. 11 \(10\): e0164272.](#)
12. Scanlon, V.M. *et al.* (2022) Multiparameter analysis of timelapse imaging reveals kinetics of megakaryocytic erythroid progenitor clonal expansion and differentiation. [Sci Rep. 12 \(1\): 16218.](#)

<b>Storage</b>	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted.
<b>Guarantee</b>	12 months from date of despatch
<b>Acknowledgements</b>	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts
<b>Health And Safety Information</b>	Material Safety Datasheet documentation #20471 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA506SBB700">https://www.bio-rad-antibodies.com/SDS/MCA506SBB700</a>
<b>Regulatory</b>	For research purposes only

## Related Products

### Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

**Product inquiries:** [www.bio-rad-antibodies.com/technical-support](http://www.bio-rad-antibodies.com/technical-support)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](http://bio-rad-antibodies.com/datasheets)

'M421944:230815'

**Printed on 23 Jan 2026**